

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> <b>C12N 15/12, 15/24, C07K 14/725, 14/54, A61K 38/17, 38/20, C12N 5/10, C07K 16/28</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 97/04097</b> <b>(43) International Publication Date:</b> 6 February 1997 (06.02.97)
<b>(21) International Application Number:</b> PCT/US96/11889 <b>(22) International Filing Date:</b> 18 July 1996 (18.07.96)  <b>(30) Priority Data:</b> 08/504,032 19 July 1995 (19.07.95) US 08/514,014 11 August 1995 (11.08.95) US  <b>(71) Applicant:</b> GENETICS INSTITUTE, INC. [US/US]; 87 Cambridge Park Drive, Cambridge, MA 02140 (US).  <b>(72) Inventors:</b> JACOBS, Kenneth; 151 Beaumont Avenue, Newton, MA 02160 (US). KELLEHER, Kerry; 50 Hurley Circle, Marlborough, MA 01752 (US). CARLIN, McKeough; 16 Chauncy Street #2, Cambridge, MA 02138 (US). GOLDMAN, Samuel; 9 Mohawk Drive, Acton, MA 01720 (US). PITTMAN, Debra; 20 N. Shore Road, Windham, NH 03087 (US). MI, Sha; 4 Vernon Road, Belmont, MA 02178 (US). NEBEN, Steven; 13 Duggan Road, Acton, MA 01720 (US). GIANNOTTI, JoAnn; 409 Arlington Street, Acton, MA 01720 (US). GOLDEN-FLEET, Margaret; 19 Usher Road, Medford, MA 02155 (US).  <b>(74) Agent:</b> BROWN, Scott, A.; Genetics Institute, Inc., 87 CambridgePark Drive, Cambridge, MA 02140 (US).	<b>(81) Designated States:</b> AU, CA, JP, MX, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>	
<b>(54) Title:</b> HUMAN CTLA-8 AND USES OF CTLA-8-RELATED PROTEINS  <b>(57) Abstract</b>  Polynucleotides encoding human CTLA-8 and related proteins are disclosed. Human CTLA-8 proteins and methods for their production are also disclosed. Methods of treatment using human CTLA-8 proteins, rat CTLA-8 proteins and herpesvirus herpes CTLA-8 proteins are also provided.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

## HUMAN CTLA-8 AND USES OF CTLA-8-RELATED PROTEINS

This application is a continuation-in-part of application Ser. No. 08/504,032, filed July 19, 1995, and a continuation-in-part of application Ser. No. 08/514,014, filed August 11, 1995.

Field of the Invention

5           The present invention relates to human CTLA-8 proteins, nucleic acids encoding such proteins, methods of treatment using such proteins. The invention also relates to the use of rat CTLA-8 proteins and herpesvirus *Saimiri* ORF13 proteins in methods of treatment.

10       Background of the Invention

          Cytokines are secreted proteins which act on specific hematopoietic target cells to cause a differentiation event or on other target cells to induce a particular physiological response, such as secretion of proteins characteristic of inflammation. Cytokines, also variously known as lymphokines, hematopoietins, interleukins, colony  
15       stimulating factors, and the like, can be important therapeutic agents, especially for diseases or conditions in which a specific cell population is depleted. For example, erythropoietin, G-CSF, and GM-CSF, have all become important for treatment of anemia and leukopenia, respectively. Other cytokines such as interleukin-3, interleukin-6, interleukin-11 and interleukin-12 show promise in treatment of  
20       conditions such as thrombocytopenia and modulation of immune response.

          For these reasons a significant research effort has been expended in searching for novel cytokines and cloning the DNAs which encode them. In the past, novel cytokines were identified by assaying a particular cell such as a bone marrow cell, for a measurable response, such as proliferation. The search for novel cytokines has thus  
5       been limited by the assays available, and if a novel cytokine has an activity which is unmeasurable by a known assay, the cytokine remains undetectable. In a newer approach, cDNAs encoding cytokines have been detected using the polymerase chain

reaction (PCR) and oligonucleotide primers having homology to shared motifs of known cytokines or their receptors. The PCR approach is also limited by the necessity for knowledge of previously cloned cytokines in the same protein family. Cytokines have also been cloned using subtractive hybridization to construct and screen cDNA libraries, or they can potentially be cloned using PCR followed by gel electrophoresis to detect differentially expressed genes. The subtractive hybridization methods are based on the assumption that cytokine mRNAs are those that are differentially expressed, and these methods do not require any prior knowledge of the sequence of interest. However, many cytokines may be encoded by mRNAs which are not differentially expressed, and thus are undetectable using these methods.

It would be desirable to develop new methods for identifying novel cytokines and other secreted factors and to isolate polynucleotides encoding them.

#### Summary of the Invention

In developing the present invention, methods were employed which selectively identify polynucleotides which encode secreted proteins. One such polynucleotide was isolated which encodes "human CTLA-8." In accordance with the present invention, polynucleotides encoding human CTLA-8 and active fragments thereof are disclosed. "CTLA-8" is used throughout the present specification to refer to both proteins and polynucleotides encoding those proteins and to refer to proteins and polynucleotides from all mammalian species.

In certain embodiments, the present invention provides an isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:1 from nucleotide 146 to nucleotide 544;
- (b) a nucleotide sequence capable of hybridizing to a nucleic acid sequence specified in (a);
- (c) a nucleotide sequence varying from the sequence of the nucleotide sequence specified in (a) as a result of degeneracy of the genetic code; and
- (d) an allelic variant of the nucleotide sequence specified in (a).



Preferably, the polynucleotide of the invention encodes a protein having CTLA-8 activity. In other embodiments the polynucleotide is operably linked to an expression control sequence. In other preferred embodiments, the polynucleotide is contained in a vector suitable for *in vivo* expression in a mammalian subject. Polynucleotides comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 55 to nucleotide 544, the nucleotide sequence of SEQ ID NO:1 from nucleotide 139 to nucleotide 544 or the nucleotide sequence of SEQ ID NO:1 from nucleotide 86 to nucleotide 544 are particularly preferred.

Host cells transformed with the polynucleotides of the invention are also provided, including mammalian cells.

Processes are also provided for producing a human CTLA-8 protein, said processes comprising:

- (a) growing a culture of the host cell of the invention in a suitable culture medium; and
- (b) purifying the human CTLA-8 protein from the culture.

Isolated human CTLA-8 protein is also provided which comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:2;
  - (b) the amino acid sequence of SEQ ID NO:2 from amino acids 11 to 163;
  - (c) the amino acid sequence of SEQ ID NO:2 from amino acids 29 to 163;
  - (d) the amino acid sequence of SEQ ID NO:2 from amino acids 31 to 163;
- and
- (e) fragments of (a), (b), (c) or (d) having CTLA-8 activity.

Proteins comprising the amino acid sequence of SEQ ID NO:2 and comprising the sequence from amino acids 29 to 163, from amino acid 31 to 163, or from amino acids 11 to 163 of SEQ ID NO:2 are particularly preferred. Preferably, the protein has CTLA-8 activity. Pharmaceuticals composition comprising a human CTLA-8 protein of the invention and a pharmaceutically acceptable carrier are also provided.

Compositions are also disclosed which comprise an antibody which specifically reacts with a human CTLA-8 protein of the invention.

Methods of treating a mammalian subject are also provided which comprise administering a therapeutically effective amount of a pharmaceutical composition comprising a human CTLA-8 protein.

5 Rat CTLA-8 and active (*i.e.*, having CTLA-8 activity) fragments thereof may also be used in such methods of treatment. Preferably the rat protein is administered as a composition comprising a pharmaceutically acceptable carrier and a protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:4;
- (b) the amino acid sequence of SEQ ID NO:4 from amino acids 18 to 150;
- 10 and
- (c) fragments of (a) or (b) having CTLA-8 activity.

Herpesvirus *Saimiri* ORF13, referred to herein as "herpes CTLA-8", and active (*i.e.*, having CTLA-8 activity) fragments thereof and active fragments thereof may also be used in such methods of treatment. Preferably the herpes CTLA-8 protein is  
15 administered as a composition comprising a pharmaceutically acceptable carrier and a protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:6;
- (b) the amino acid sequence of SEQ ID NO:6 from amino acids 19 to 151;
- 20 and
- (c) fragments of (a) or (b) having CTLA-8 activity.

The invention also provides a method of treating a mammalian subject comprising administering a therapeutically effective amount of a composition comprising a pharmaceutically acceptable carrier and IL-17 or an active fragment thereof.

25 In methods of treatment provided by the present invention, preferably the subject is treated to produce an effect selected from the group consisting of inhibition of angiogenesis, inhibition of growth or proliferation of vascular endothelial cells, inhibition of tumor growth, inhibition of angiogenesis-dependent tissue growth, proliferation of myeloid cells or progenitors, proliferation of erythroid cells or  
30 progenitors, proliferation of lymphoid cells or progenitors, induction of IFN $\gamma$  production, induction of IL-3 production and induction of GM-CSF production.

### Brief Description of the Figures

Fig. 1 is a comparison of homologous regions of the amino acid sequences of human CTLA-8 (indicated as "B18\_F1"), rat CTLA-8 (indicated as "Musctla8") and herpes CTLA-8 (indicated as "Hsvie\_2").

5 Fig. 2 depicts autoradiographs demonstrating expression of human CTLA-8 in COS cells.

Fig. 3 presents data relating to the ability of human CTLA-8 to inhibit angiogenesis.

10 Figs. 4 and 5 present data relating to the ability of human CTLA-8 to produce or induce hematopoietic activity.

Figs. 6 and 7 present data demonstrating the ability of human CTLA-8 to induce production of IL-6 and IL-8.

### Detailed Description of Preferred Embodiments

15 The inventors of the present application have identified and provided a polynucleotide encoding a human CTLA-8 protein. SEQ ID NO:1 provides the nucleotide sequence of a cDNA encoding the human CTLA-8 protein. SEQ ID NO:2 provides the amino acid sequence of the human CTLA-8 protein. Alternatively, the initiating methionine may be at amino acid 11 of SEQ ID NO:2. On the basis of  
20 amino terminal sequencing, the mature protein sequence is believed to begin at amino acid 31 of SEQ ID NO:2 (encoded by the sequence beginning with nucleotide 146 of SEQ ID NO:1).

The region from amino acid 29 to amino acid 163 of human CTLA-8 (SEQ ID NO:2) shows marked homology to portions of rat CTLA-8 (amino acids 18 to 150 of  
25 SEQ ID NO:4) and herpesvirus *Saimiri* ORF13 ("herpes CTLA-8") (amino acids 19 to 151 of SEQ ID NO:5). A cDNA sequence encoding rat CTLA-8 is listed at SEQ ID NO:3 and its corresponding amino acid sequence is reported at SEQ ID NO:4. A cDNA sequence encoding herpes CTLA-8 is listed at SEQ ID NO:5 and its corresponding amino acid sequence is reported at SEQ ID NO:6. Homology between  
30 rat CTLA-8 and herpes CTLA-8 was reported by Rouvier et al., J. Immunol. 1993, 150, 5445-5456.

Applicants had previously incorrectly identified the rat sequences of SEQ ID NO:3 and SEQ ID NO:4 as applying to murine CTLA-8. Applicants' human CTLA-8 (B18) does also show homolgy to the true murine CTLA-8 sequence.

5       Golstein et al. (WO95/18826; Fossiez et al., Microbial Evasion and Subversion of Immunity 544:3222 (Abstract)) have also reported a species they initially identified as "human CTLA-8." However, examination of the sequence of the Golstein et al. species and the human CTLA-8 (B18) sequence of the present invention readily reveals that they are two different proteins, although they are homologous with each other and with the rat CTLA-8 and herpes CTLA-8 identified herein. The Golstein  
10       et al. species has now been renamed as interleukin-17 (IL-17). Because of the homology between applicants' human CTLA-8 (B18) and IL-17, these proteins are expected to share some activities.

It has also been preliminarily determined that human CTLA-8 (B18) forms homodimers when expressed. As a result, human CTLA-8 proteins may possess  
15       activity in either monomeric or dimeric forms. Human CTLA-8 proteins can also be produced as heterodimers with rat and herpes CTLA-8 proteins and with human IL-17. These heterodimers are also expected to have activities of the proteins of which they are comprised.

Forms of human CTLA-8 protein of less than full length are encompassed  
20       within the present invention and may be produced by expressing a corresponding fragment of the polynucleotide encoding the human CTLA-8 protein (SEQ ID NO:1). These corresponding polynucleotide fragments are also part of the present invention. Modified polynucleotides as described above may be made by standard molecular biology techniques, including site-directed mutagenesis methods which are known in  
25       the art or by the polymerase chain reaction using appropriate oligonucleotide primers.

For the purposes of the present invention, a protein has "CTLA-8 activity" if it either (1) displays biological activity in a factor-dependent cell proliferation assay (preferably an assay in which full-length the corresponding species full-length CTLA-8 is active) (including without limitation those assays described below), or (2) induces  
30       expression or secretion of  $\gamma$ -IFN, or (3) displays chemoattractant of chemotactic activity in a chemoattraction or chemotaxis assay (preferably as assay in which full-

length the corresponding species full-length CTLA-8 is active) or (4) induces expression of secretion of IL-3 or GM-CSF.

Human CTLA-8 protein or fragments thereof having CTLA-8 activity may be fused to carrier molecules such as immunoglobulins. For example, human CTLA-8 protein may be fused through "linker" sequences to the Fc portion of an immunoglobulin.

The invention also encompasses allelic variations of the nucleotide sequence as set forth in SEQ ID NO:1, that is, naturally-occurring alternative forms of the isolated polynucleotide of SEQ ID NO:1 which also encode human CTLA-8 or CTLA-8 proteins having CTLA-8 activity. Also included in the invention are isolated polynucleotides which hybridize to the nucleotide sequence set forth in SEQ ID NO:1 under highly stringent (0.2xSSC at 65°C), stringent (e.g. 4xSSC at 65°C or 50% formamide and 4xSSC at 42°C), or relaxed (4xSSC at 50°C or 30-40% formamide and 4xSSC at 42°C) conditions. Isolated polynucleotides which encode human CTLA-8 protein but which differ from the nucleotide sequence set forth in SEQ ID NO:1 by virtue of the degeneracy of the genetic code are also encompassed by the present invention. Variations in the nucleotide sequence as set forth in SEQ ID NO:1 which are caused by point mutations or by induced modifications which enhance CTLA-8 activity, half-life or production level are also included in the invention.

The isolated polynucleotides of the invention may be operably linked to an expression control sequence such as the pMT2 or pED expression vectors disclosed in Kaufman et al., Nucleic Acids Res. 19, 4485-4490 (1991), in order to produce the CTLA-8 protein recombinantly. Many suitable expression control sequences are known in the art. General methods of expressing recombinant proteins are also known and are exemplified in R. Kaufman, Methods in Enzymology 185, 537-566 (1990). As defined herein "operably linked" means enzymatically or chemically ligated to form a covalent bond between the isolated polynucleotide of the invention and the expression control sequence, in such a way that the CTLA-8 protein is expressed by a host cell which has been transformed (transfected) with the ligated polynucleotide/expression control sequence.

A number of types of cells may act as suitable host cells for expression of the human CTLA-8 protein. Any cell type capable of expressing functional human CTLA-8 protein may be used. Suitable mammalian host cells include, for example, monkey COS cells, Chinese Hamster Ovary (CHO) cells, human kidney 293 cells, human epidermal A431 cells, human Colo205 cells, 3T3 cells, CV-1 cells, other transformed primate cell lines, normal diploid cells, cell strains derived from in vitro culture of primary tissue, primary explants, HeLa cells, mouse L cells, BHK, HL-60, U937, HaK, Rat2, BaF3, 32D, FDCP-1, PC12 or C2C12 cells.

The human CTLA-8 protein may also be produced by operably linking the isolated polynucleotide of the invention to suitable control sequences in one or more insect expression vectors, and employing an insect expression system. Materials and methods for baculovirus/insect cell expression systems are commercially available in kit form from, e.g., Invitrogen, San Diego, California, U.S.A. (the MaxBac® kit), and such methods are well known in the art, as described in Summers and Smith, Texas Agricultural Experiment Station Bulletin No. 1555 (1987), incorporated herein by reference. Soluble forms of the human CTLA-8 protein may also be produced in insect cells using appropriate isolated polynucleotides as described above.

Alternatively, the human CTLA-8 protein may be produced in lower eukaryotes such as yeast or in prokaryotes such as bacteria. Suitable yeast strains include *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces* strains, *Candida*, or any yeast strain capable of expressing heterologous proteins. Suitable bacterial strains include *Escherichia coli*, *Bacillus subtilis*, *Salmonella typhimurium*, or any bacterial strain capable of expressing heterologous proteins.

The human CTLA-8 protein of the invention may also be expressed as a product of transgenic animals, e.g., as a component of the milk of transgenic cows, goats, pigs, or sheep which are characterized by somatic or germ cells containing a polynucleotide sequence encoding the human CTLA-8 protein.

The human CTLA-8 protein of the invention may be prepared by growing a culture of transformed host cells under culture conditions necessary to express the desired protein. The resulting expressed protein may then be purified from the culture medium or cell extracts. Soluble forms of the human CTLA-8 protein of the invention

can be purified from conditioned media. Membrane-bound forms of human CTLA-8 protein of the invention can be purified by preparing a total membrane fraction from the expressing cell and extracting the membranes with a non-ionic detergent such as Triton X-100.

5           The human CTLA-8 protein can be purified using methods known to those skilled in the art. For example, the human CTLA-8 protein of the invention can be concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. Following the concentration step, the concentrate can be applied to a purification matrix such as a gel filtration medium.

10           Alternatively, an anion exchange resin can be employed, for example, a matrix or substrate having pendant diethylaminoethyl (DEAE) groups. The matrices can be acrylamide, agarose, dextran, cellulose or other types commonly employed in protein purification. Alternatively, a cation exchange step can be employed. Suitable cation exchangers include various insoluble matrices comprising sulfopropyl or

15           carboxymethyl groups. Sulfopropyl groups are preferred (e.g., S-Sepharose® columns). The purification of the human CTLA-8 protein from culture supernatant may also include one or more column steps over such affinity resins as concanavalin A-agarose, heparin-toyopearl® or Cibacrom blue 3GA Sepharose®; or by hydrophobic interaction chromatography using such resins as phenyl ether, butyl ether,

20           or propyl ether; or by immunoaffinity chromatography. Finally, one or more reverse-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, e.g., silica gel having pendant methyl or other aliphatic groups, can be employed to further purify the human CTLA-8 protein. Some or all of the foregoing purification steps, in various combinations or with other known

25           methods, can also be employed to provide a substantially purified isolated recombinant protein.

          Preferably, the human CTLA-8 protein is purified so that it is substantially free of other mammalian proteins.

          It is believed that human CTLA-8, active fragments and variants thereof, and

30           CTLA-8 related proteins (such as, for example, rat CTLA-8 and herpes CTLA-8) (collectively "CTLA-8 proteins") possess or induce cytokine activities. Human

CTLA-8 expression correlated with  $\gamma$ -IFN expression in induced primary cells and can induce the expression of IL-3 and/or GM-CSF, which expression can in turn produce effects associated with the induced cytokine. Therefore, human CTLA-8 and CTLA-8 related proteins may have an effect on proliferation or function of myeloid cells, erythroid cells, lymphoid cells and their progenitors. Human CTLA-8 proteins may also play a role in formation of platelets or their progenitors.

A protein of the present invention may exhibit cytokine, cell proliferation (either inducing or inhibiting) or cell differentiation (either inducing or inhibiting) activity or may induce production of other cytokines in certain cell populations. Many protein factors discovered to date, including all known cytokines, have exhibited activity in one or more factor dependent cell proliferation assays, and hence the assays serve as a convenient confirmation of cytokine activity. The activity of a protein of the present invention is evidenced by any one of a number of routine factor dependent cell proliferation assays for cell lines including, without limitation, 32D, DA2, DA1G, T10, B9, B9/11, BaF3, MC9/G, M+ (preB M+), 2E8, RB5, DA1, 123, T1165, HT2, CTLL2, TF-1, Mo7e and CMK.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for T-cell or thymocyte proliferation include without limitation those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, *In Vitro* assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, *Immunologic studies in Humans*); Takai et al., *J. Immunol.* 137:3494-3500, 1986; Bertagnolli et al., *J. Immunol.* 145:1706-1712, 1990; Bertagnolli et al., *Cellular Immunology* 133:327-341, 1991; Bertagnolli, et al., *J. Immunol.* 149:3778-3783, 1992; Bowman et al., *J. Immunol.* 152: 1756-1761, 1994.

Assays for cytokine production and/or proliferation of spleen cells, lymph node cells or thymocytes include, without limitation, those described in: *Polyclonal T cell stimulation*, Kruisbeek, A.M. and Shevach, E.M. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 3.12.1-3.12.14, John Wiley and Sons, Toronto. 1994; and *Measurement of mouse and human Interferon  $\gamma$* , Schreiber, R.D.



In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.8.1-6.8.8, John Wiley and Sons, Toronto. 1994.

Assays for proliferation and differentiation of hematopoietic and lymphopoietic cells include, without limitation, those described in: Measurement of Human and Murine Interleukin 2 and Interleukin 4, Bottomly, K., Davis, L.S. and Lipsky, P.E. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.3.1-6.3.12, John Wiley and Sons, Toronto. 1991; deVries et al., J. Exp. Med. 173:1205-1211, 1991; Moreau et al., Nature 336:690-692, 1988; Greenberger et al., Proc. Natl. Acad. Sci. U.S.A. 80:2931-2938, 1983; Measurement of mouse and human interleukin 6 - Nordan, R. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.6.1-6.6.5, John Wiley and Sons, Toronto. 1991; Smith et al., Proc. Natl. Acad. Sci. U.S.A. 83:1857-1861, 1986; Measurement of human Interleukin 11 - Bennett, F., Giannotti, J., Clark, S.C. and Turner, K. J. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.15.1 John Wiley and Sons, Toronto. 1991; Measurement of mouse and human Interleukin 9 - Ciarletta, A., Giannotti, J., Clark, S.C. and Turner, K.J. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.13.1, John Wiley and Sons, Toronto. 1991.

Assays for T-cell clone responses to antigens (which will identify, among others, proteins that affect APC-T cell interactions as well as direct T-cell effects by measuring proliferation and cytokine production) include, without limitation, those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober  
Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function; Chapter 6, Cytokines and their cellular receptors; Chapter 7, Immunologic studies in Humans); Weinberger et al., Proc. Natl. Acad. Sci. USA 77:6091-6095, 1980; Weinberger et al., Eur. J. Immun. 11:405-411, 1981; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988.

A protein of the present invention may also exhibit immune stimulating or immune suppressing activity, including without limitation the activities for which assays are described herein. A protein may be useful in the treatment of various

immune deficiencies and disorders (including severe combined immunodeficiency (SCID)), e.g., in regulating (up or down) growth and proliferation of T and/or B lymphocytes, as well as effecting the cytolytic activity of NK cells and other cell populations. These immune deficiencies may be genetic or be caused by viral (e.g.,  
5 HIV) as well as bacterial or fungal infections, or may result from autoimmune disorders. More specifically, infectious diseases caused by viral, bacterial, fungal or other infection may be treatable using a protein of the present invention, including infections by HIV, hepatitis viruses, herpes viruses, mycobacteria, leishmania, malaria and various fungal infections such as candida. Of course, in this regard, a protein of  
10 the present invention may also be useful where a boost to the immune system generally would be indicated, i.e., in the treatment of cancer.

Autoimmune disorders which may be treated using a protein of the present invention include, for example, multiple sclerosis, systemic lupus erythematosus, rheumatoid arthritis, autoimmune pulmonary inflammation, Guillain-Barre syndrome,  
15 autoimmune thyroiditis, insulin dependent diabetes mellitus, myasthenia gravis, graft-versus-host disease and autoimmune inflammatory eye disease. Such a protein of the present invention may also be useful in the treatment of allergic reactions and conditions, such as asthma or other respiratory problems. Other conditions, in which immune suppression is desired (including, for example, asthma and related respiratory  
20 conditions), may also be treatable using a protein of the present invention.

A protein of the present invention may also suppress chronic or acute inflammation, such as, for example, that associated with infection (such as septic shock or systemic inflammatory response syndrome (SIRS)), inflammatory bowel disease, Crohn's disease or resulting from over production of cytokines such as TNF  
25 or IL-1 (such as the effect demonstrated by IL-11).

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for thymocyte or splenocyte cytotoxicity include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing  
30 Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte

- Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al., J. Immunol. 135:1564-1572, 1985; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988;
- 5 Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al., J. Immunol. 135:1564-1572, 1985; Takai et al., J. Immunol. 137:3494-3500, 1986; Bowman et al., J. Virology 61:1992-1998; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnolli et al., Cellular Immunology 133:327-341, 1991; Brown et al., J. Immunol. 153:3079-3092, 1994.
- 10 Assays for T-cell-dependent immunoglobulin responses and isotype switching (which will identify, among others, proteins that modulate T-cell dependent antibody responses and that affect Th1/Th2 profiles) include, without limitation, those described in: Maliszewski, J. Immunol. 144:3028-3033, 1990; and Assays for B cell function: *In vitro* antibody production, Mond, J.J. and Brunswick, M. In *Current*
- 15 *Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 3.8.1-3.8.16, John Wiley and Sons, Toronto. 1994.
- Mixed lymphocyte reaction (MLR) assays (which will identify, among others, proteins that generate predominantly Th1 and CTL responses) include, without limitation, those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing
- 20 Associates and Wiley-Interscience (Chapter 3, *In Vitro* assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnolli et al., J. Immunol. 149:3778-3783, 1992.
- 25 Dendritic cell-dependent assays (which will identify, among others, proteins expressed by dendritic cells that activate naive T-cells) include, without limitation, those described in: Guery et al., J. Immunol. 134:536-544, 1995; Inaba et al., Journal of Experimental Medicine 173:549-559, 1991; Macatonia et al., Journal of Immunology 154:5071-5079, 1995; Porgador et al., Journal of Experimental Medicine
- 30 182:255-260, 1995; Nair et al., Journal of Virology 67:4062-4069, 1993; Huang et al., Science 264:961-965, 1994; Macatonia et al., Journal of Experimental Medicine

169:1255-1264, 1989; Bhardwaj et al., Journal of Clinical Investigation 94:797-807, 1994; and Inaba et al., Journal of Experimental Medicine 172:631-640, 1990.

Assays for lymphocyte survival/apoptosis (which will identify, among others, proteins that prevent apoptosis after superantigen induction and proteins that regulate lymphocyte homeostasis) include, without limitation, those described in:

5 Darzynkiewicz et al., Cytometry 13:795-808, 1992; Gorczyca et al., Leukemia 7:659-670, 1993; Gorczyca et al., Cancer Research 53:1945-1951, 1993; Itoh et al., Cell 66:233-243, 1991; Zacharchuk, Journal of Immunology 145:4037-4045, 1990; Zamai et al., Cytometry 14:891-897, 1993; Gorczyca et al., International Journal of

10 Oncology 1:639-648, 1992.

Assays for proteins that influence early steps of T-cell commitment and development include, without limitation, those described in: Antica et al., Blood 84:111-117, 1994; Fine et al., Cellular Immunology 155:111-122, 1994; Galy et al., Blood 85:2770-2778, 1995; Toki et al., Proc. Nat. Acad Sci. USA 88:7548-7551,

15 1991.

A protein of the present invention may be useful in regulation of hematopoiesis and, consequently, in the treatment of myeloid or lymphoid cell deficiencies. Even marginal biological activity in support of colony forming cells or of factor-dependent cell lines indicates involvement in regulating hematopoiesis, e.g. in supporting the

20 growth and proliferation of erythroid progenitor cells alone or in combination with other cytokines, thereby indicating utility, for example, in treating various anemias or for use in conjunction with irradiation/chemotherapy to stimulate the production of erythroid precursors and/or erythroid cells; in supporting the growth and proliferation of myeloid cells such as granulocytes and monocytes/macrophages (i.e., traditional

25 CSF activity) useful, for example, in conjunction with chemotherapy to prevent or treat consequent myelo-suppression; in supporting the growth and proliferation of megakaryocytes and consequently of platelets thereby allowing prevention or treatment of various platelet disorders such as thrombocytopenia, and generally for use in place of or complementarily to platelet transfusions; and/or in supporting the growth

30 and proliferation of hematopoietic stem cells which are capable of maturing to any and all of the above-mentioned hematopoietic cells and therefore find therapeutic utility

in various stem cell disorders (such as those usually treated with transplantation, including, without limitation, aplastic anemia and paroxysmal nocturnal hemoglobinuria), as well as in repopulating the stem cell compartment post irradiation/chemotherapy, either *in-vivo* or *ex-vivo* (i.e. in conjunction with bone marrow transplantation) as normal cells or genetically manipulated for gene therapy.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for proliferation and differentiation of various hematopoietic lines are cited above.

Assays for embryonic stem cell differentiation (which will identify, among others, proteins that influence embryonic differentiation hematopoiesis) include, without limitation, those described in: Johansson et al. *Cellular Biology* 15:141-151, 1995; Keller et al., *Molecular and Cellular Biology* 13:473-486, 1993; McClanahan et al., *Blood* 81:2903-2915, 1993.

Assays for stem cell survival and differentiation (which will identify, among others, proteins that regulate lympho-hematopoiesis) include, without limitation, those described in: Methylcellulose colony forming assays, Freshney, M.G. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 265-268, Wiley-Liss, Inc., New York, NY. 1994; Hirayama et al., *Proc. Natl. Acad. Sci. USA* 89:5907-5911, 1992; Primitive hematopoietic colony forming cells with high proliferative potential, McNiece, I.K. and Briddell, R.A. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 23-39, Wiley-Liss, Inc., New York, NY. 1994; Neben et al., *Experimental Hematology* 22:353-359, 1994; Cobblestone area forming cell assay, Ploemacher, R.E. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 1-21, Wiley-Liss, Inc., New York, NY. 1994; Long term bone marrow cultures in the presence of stromal cells, Spooncer, E., Dexter, M. and Allen, T. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 163-179, Wiley-Liss, Inc., New York, NY. 1994; Long term culture initiating cell assay, Sutherland, H.J. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 139-162, Wiley-Liss, Inc., New York, NY. 1994.

CTLA-8 proteins are useful in the treatment of various immune deficiencies and disorders (including SCID), *e.g.*, in regulating (up or down) growth, proliferation and/or activity of T and/or B lymphocytes, as well as the cytolytic activity of NK cells. These immune deficiencies may be caused by viral (*e.g.*, HIV) as well as bacterial infections, or may result from autoimmune disorders. More specifically, infectious diseases caused by viral, bacterial, fungal or other infection may be treatable using CTLA-8 proteins, including infections by HIV, hepatitis, influenza, CMV, herpes, mycobacterium, leishmaniasis, malaria and various fungal infections (such as candida). Of course, in this regard, the CTLA-8 proteins may also be useful where a boost to the immune system generally would be indicated, *i.e.*, in the treatment of cancer or as an adjuvant to vaccines. Autoimmune disorders which may be treated using factors of the present invention include, for example, multiple sclerosis, systemic lupus erythematosus, rheumatoid arthritis, autoimmune pulmonary inflammation, Guillain-Barre syndrome, autoimmune thyroiditis, insulin dependent diabetes melitis and autoimmune inflammatory eye disease. The CTLA-8 proteins are also expected to be useful in the treatment of allergic reactions and conditions.

CTLA-8 proteins are also expected to have chemotactic activity. A protein or peptide has "chemotactic activity," as used herein, if it can stimulate, directly or indirectly, the directed orientation or movement of cells, including myeloid and lymphoid cells. Preferably, the protein or peptide has the ability to directly stimulate directed movement of cells (particularly T-cells). Whether a particular protein or peptide has chemotactic activity for cells can be readily determined by employing such protein or peptide in any known assay for cell chemotaxis.

CTLA-8 proteins also inhibit growth and proliferation of vascular endothelial cells. As a result, human CTLA-8 proteins are effective in inhibiting angiogenesis (*i.e.*, vascular formation). This activity will also be useful in the treatment of tumors and other conditions in which angiogenesis is involved. Inhibition of angiogenesis by human CTLA-8 proteins will also result in inhibition or prevention of the condition to which normal angiogenesis would contribute.

Isolated CTLA-8 proteins, purified from cells or recombinantly produced, may be used as a pharmaceutical composition when combined with a pharmaceutically

acceptable carrier. Such a composition may contain, in addition to CTLA-8 protein and carrier, diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredient(s). The characteristics of the carrier will depend on the route of administration. The pharmaceutical composition of the invention may also contain cytokines, lymphokines, or other hematopoietic factors such as M-CSF, GM-CSF, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-15, G-CSF,  $\gamma$ -IFN, stem cell factor, and erythropoietin. The pharmaceutical composition may contain thrombolytic or anti-thrombotic factors such as plasminogen activator and Factor VIII. The pharmaceutical composition may further contain other anti-inflammatory agents. Such additional factors and/or agents may be included in the pharmaceutical composition to produce a synergistic effect with CTLA-8 protein, or to minimize side effects caused by the CTLA-8 protein. Conversely, CTLA-8 protein may be included in formulations of the particular cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent to minimize side effects of the cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent.

The pharmaceutical composition of the invention may be in the form of a liposome in which CTLA-8 protein is combined, in addition to other pharmaceutically acceptable carriers, with amphipathic agents such as lipids which exist in aggregated form as micelles, insoluble monolayers, liquid crystals, or lamellar layers which in aqueous solution. Suitable lipids for liposomal formulation include, without limitation, monoglycerides, diglycerides, sulfatides, lysolecithin, phospholipids, saponin, bile acids, and the like. Preparation of such liposomal formulations is within the level of skill in the art, as disclosed, for example, in U.S. Patent No. 4,235,871; U.S. Patent No. 4,501,728; U.S. Patent No. 4,837,028; and U.S. Patent No. 4,737,323, all of which are incorporated herein by reference.

As used herein, the term "therapeutically effective amount" means the total amount of each active component of the pharmaceutical composition or method that is sufficient to show a meaningful patient benefit, e.g., amelioration of symptoms of,

healing of, or increase in rate of healing of such conditions. When applied to an individual active ingredient, administered alone, the term refers to that ingredient alone. When applied to a combination, the term refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in  
5 combination, serially or simultaneously.

In practicing the method of treatment or use of the present invention, a therapeutically effective amount of CTLA-8 protein is administered to a mammal. CTLA-8 protein may be administered in accordance with the method of the invention either alone or in combination with other therapies such as treatments employing  
10 cytokines, lymphokines or other hematopoietic factors. When co-administered with one or more cytokines, lymphokines, other hematopoietic factors or vaccine components (such as antigens or other adjuvants), CTLA-8 protein may be administered either simultaneously with the cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors, or sequentially. If  
15 administered sequentially, the attending physician will decide on the appropriate sequence of administering CTLA-8 protein in combination with cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors.

Administration of CTLA-8 protein used in the pharmaceutical composition or to practice the method of the present invention can be carried out in a variety of  
20 conventional ways, such as oral ingestion, inhalation, or cutaneous, subcutaneous, or intravenous injection. Intravenous administration to the patient is preferred.

When a therapeutically effective amount of CTLA-8 protein is administered orally, CTLA-8 protein will be in the form of a tablet, capsule, powder, solution or elixir. When administered in tablet form, the pharmaceutical composition of the  
25 invention may additionally contain a solid carrier such as a gelatin or an adjuvant. The tablet, capsule, and powder contain from about 5 to 95% CTLA-8 protein, and preferably from about 25 to 90% CTLA-8 protein. When administered in liquid form, a liquid carrier such as water, petroleum, oils of animal or plant origin such as peanut oil, mineral oil, soybean oil, or sesame oil, or synthetic oils may be added. The liquid  
30 form of the pharmaceutical composition may further contain physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol,



propylene glycol or polyethylene glycol. When administered in liquid form, the pharmaceutical composition contains from about 0.5 to 90% by weight of CTLA-8 protein, and preferably from about 1 to 50% CTLA-8 protein.

5 When a therapeutically effective amount of CTLA-8 protein is administered by intravenous, cutaneous or subcutaneous injection, CTLA-8 protein will be in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable protein solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred pharmaceutical composition for intravenous, cutaneous, or subcutaneous injection should contain, in  
10 addition to CTLA-8 protein an isotonic vehicle such as Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, Lactated Ringer's Injection, or other vehicle as known in the art. The pharmaceutical composition of the present invention may also contain stabilizers, preservatives, buffers, antioxidants, or other additive known to those of skill in the art.

15 The amount of CTLA-8 protein in the pharmaceutical composition of the present invention will depend upon the nature and severity of the condition being treated, and on the nature of prior treatments which the patient has undergone. Ultimately, the attending physician will decide the amount of CTLA-8 protein with which to treat each individual patient. Initially, the attending physician will  
20 administer low doses of CTLA-8 protein and observe the patient's response. Larger doses of CTLA-8 protein may be administered until the optimal therapeutic effect is obtained for the patient, and at that point the dosage is not generally increased further. It is contemplated that the various pharmaceutical compositions used to practice the method of the present invention should contain about 0.1  $\mu$ g to about 100 mg of  
25 CTLA-8 protein per kg body weight, preferably about 0.1  $\mu$ g to about 10 mg of CTLA-8 protein per kg body weight, more preferably about 0.1  $\mu$ g to about 100  $\mu$ g of CTLA-8 protein per kg body weight, most preferably preferably about 0.1  $\mu$ g to about 10  $\mu$ g of CTLA-8 protein per kg body weight.

30 The duration of intravenous therapy using the pharmaceutical composition of the present invention will vary, depending on the severity of the disease being treated and the condition and potential idiosyncratic response of each individual patient. It

is contemplated that the duration of each application of the CTLA-8 protein will be in the range of 12 to 24 hours of continuous intravenous administration. Ultimately the attending physician will decide on the appropriate duration of intravenous therapy using the pharmaceutical composition of the present invention.

5 CTLA-8 protein of the invention may also be used to immunize animals to obtain polyclonal and monoclonal antibodies which specifically react with the CTLA-8 protein and which may inhibit CTLA-8 binding to its receptor. Such antibodies are also useful for performing diagnostics assays for CTLA-8 in accordance with known methods. Such antibodies may be obtained using the entire CTLA-8 protein as an  
10 immunogen, or by using fragments of human CTLA-8 protein. The peptide immunogens additionally may contain a cysteine residue at the carboxyl terminus, and are conjugated to a hapten such as keyhole limpet hemocyanin (KLH). Additional peptide immunogens may be generated by replacing tyrosine residues with sulfated tyrosine residues. Methods for synthesizing such peptides are known in the art, for  
15 example, as in R.P. Merrifield, J.Amer.Chem.Soc. 85, 2149-2154 (1963); J.L. Krstenansky, et al., FEBS Lett. 211, 10 (1987).

Neutralizing or non-neutralizing antibodies (preferably monoclonal antibodies) binding to human CTLA-8 protein may also be useful therapeutics for certain tumors and also in the treatment of conditions described above. These neutralizing  
20 monoclonal antibodies are capable of blocking the ligand binding to the human CTLA-8 protein or mayh promote clearance of protein from the patient.

Because of their homology to human CTLA-8, rat CTLA-8 proteins, herpes CTLA-8 proteins and IL-17 proteins (the "human CTLA-8" of Golstein et al., supra) will also possess CTLA-8 activity as described above. As a result, rat and herpes  
25 CTLA-8 proteins and IL-17 proteins, as well as active fragments and variants thereof, can be used in preparation of pharmaceutical compositions and in methods of treatment as described for human CTLA-8. Rat and herpes CTLA-8 proteins, and active fragments and variants thereof, can be produced as described above using the polynucleotides (or fragments or variants thereof) described in SEQ ID NO:3 and SEQ  
30 ID NO:5, respectively. Rat and herpes CTLA-8 may also be produced as described in Rouvier et al., J. Immunol. 1993, 150, 5445-5456. CTLA-8 proteins of other

species can also be used as described herein. cDNAs encoding rat CTLA-8 and herpes CTLA-8 were deposited with the American Type Culture Collection on July 6, 1995 and assigned accession numbers ATCC 69867 and ATCC 69866, respectively. IL-17 proteins may also be produced as described in Golstein et al., supra.

5        Because of its homology to IL-17, the human CTLA-8 (B18) proteins of the present invention may also share some activities with IL-17.

For the purposes of treatment or therapy, any of the proteins discussed or disclosed herein may be administered by *in vivo* expression of the protein in a mammalian subject. In such instances, a polynucleotide encoding the desired protein  
10        is administered to the subject in manner allowing expression in accordance with known methods, including without limitation the adenovirus methods disclosed herein.

#### Example 1

##### 15        Isolation of Human CTLA-8 cDNA

A partial clone for human CTLA-8 was isolated from a cDNA library made from RNA isolated from stimulated human peripheral blood mononuclear cells. This partial was identified as "B18." B18 is sometimes used herein to refer to the human CTLA-8 of the present invention. Homology searches identified this partial clone as  
20        being related to the herpes and rat CTLA-8 genes. DNA sequence of this partial clone was used to isolate the full-length clone.

In order to isolate a full-length cDNA for B18, a directional, full-length cDNA library by standard means in the COS expression vector pMV2. The cDNA library was transformed into *E. coli* by electroporation. The bulk of the original transformed  
25        cDNA library was frozen in glycerol at -80°C. An aliquot was titered to measure the concentration of transformed *E. coli*. The *E. coli* were thawed, diluted to 76,000/0.1 ml in media containing ampicillin, and 0.1 ml was distributed into the wells of a microtiter dish in an 8 x 8 array. The microtiter dish was placed at 37°C overnight to grow the *E. coli*.

30        To prepare DNA for PCR, 20 µl aliquots of culture from each well were withdrawn and pooled separately for each row and column of eight wells, giving 16

pools of 160  $\mu$ l each. The *E. coli* were pelleted, resuspended in 160  $\mu$ l of standard lysis buffer consisting of 10 mM TrisHCl pH8, 1 mM EDTA, 0.01% Triton X-100, and lysed by heating to 95°C for 10 minutes.

5 To identify which of the wells contained *E. coli* transformed with B18, PCR was performed first on the DNA preps corresponding to the eight columns. The PCR consisted of two sequential reactions with nested oligonucleotides using standard conditions. The oligonucleotides used for the PCR reaction were derived from the sequence of the partial B18 clone. They were:

10 B185: CACAGGCATACACAGGAAGATACATTCA (SEQ ID NO:7)  
B183: TCTTGCTGGATGGGAACGGAATTCA (SEQ ID NO:8)  
B18N: ATACATTCACAGAAGAGCTTCCTGCACA (SEQ ID NO:9)

15 The PCR conditions were 2.5 mM MgCl<sub>2</sub> and 95°C x 2 min for one cycle, 95°C x 1 min plus 68°C x 1 min for 30 cycles, and 68°C x 10 for one cycle. Each reaction was 20  $\mu$ l. The first reaction contained oligonucleotides B185 and B183 and 1  $\mu$ l of the DNA preparations. The second reaction contained oligonucleotides B183 and B18N and 1  $\mu$ l of the first reaction.

20 DNA preps that potentially contained a full-length B18 cDNA clone were identified by agarose gel electrophoresis on an aliquot of the second PCR reaction. A DNA band of the correct mobility was assumed to be derived from a B18 cDNA. Next the same sequence of PCR reactions and gel analysis was done on the DNA preps corresponding to the eight rows. The intersection of a row and a column identified well A2 as potentially containing B18, narrowing it down to the 76,000 *E.*  
25 *coli* originally seeded into that well.

To further purify the individual *E. coli* containing the putative full-length B18 cDNA clone, the concentration of *E. coli* in well A2 was measured by titring and plating dilutions of the well. Then 7600 *E. coli* were seeded into the wells of a second microtiter plate in an 8 x 8 array. The *E. coli* were grown overnight; wells were  
30 pooled, and DNA was prepared as described above. To identify which of these wells contained *E. coli* transformed with B18, sequential PCR reactions were performed

essentially as described above. Agarose gel electrophoresis identified well B2 as potentially containing a B18 cDNA.

5 The *E. coli* containing this cDNA was further purified by seeding wells of a microtiter plate with 253 *E. coli* per well and proceeding as for the purification of the *E. coli* in well A2. Well C3 was identified as containing a putative full-length B18 cDNA clone. The exact *E. coli* was identified by plating the contents of the well onto bacterial culture media and then screening the *E. coli* colonies following established protocols. The probe for these hybridizations was a PCR fragment generated by doing a PCR reaction on the B18 clone using as primers the oligonucleotides described  
10 above (SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9). When a single colony was identified, DNA was prepared and sequenced by standard methods. Comparison of this sequence to the sequence of the original partial clone confirmed identity and that the isolated cDNA was full-length.

15 The full-length clone was deposited with the American Type Culture Collection on July 6, 1995 and assigned accession number ATCC 69868.

### Example 2

#### Expression of Human CTLA-8

20 The full-length B18 clone for human CTLA-8 was transfected into COS cells which were then labelled with <sup>35</sup>S-methionine. An aliquot of conditioned medium from the transfected cell culture was reduced, denatured and electrophoresed on polyacrylamide gels. Autoradiographs of those gels are reproduced in Fig. 2. The band indicated by the arrow demonstrates expression of human CTLA-8.

### Example 3

#### Inhibition of Angiogenesis by Human CTLA-8

25 The ability of human CTLA-8 to inhibit angiogenesis was examined in an angiostatic activity assay (endothelial cell proliferation assay). The assay was done in a 96 well plate. Primary human umbilical cells (HUVECs) were seeded to 2x10<sup>3</sup>  
30 cells per well in EGM medium (Clonetics)/20% FCS and incubated at 37°C for 24 hr. The cells were then starved in M199 medium (GIBCO BRL) containing 10% charcoal

treated serum (M199-CS) for 48 hr at 37°C. Conditioned media containing B18 (human CTLA-8) was obtained from transfected COS or stably expressing CHO cells and 1:10, 1:50, 1:250, and 1:1250 dilutions prepared in M199-CS medium containing 100 ng/ml FGF. The dilutions of B18 were added to the starved cells and incubated  
5 for 72 hr at 37°C. The cells were then radiolabeled by [<sup>3</sup>H]-thymidine for 6 hr. Radiolabeled cells were washed with PBS and trypsinized for liquid scintillation counting. Results were plotted using Kaleidograph software. The results are shown in Fig. 3. In the figure, "Med" is the mock control, "B18" and "B18-1" were conditioned medium from two independent transfections of COS with DNA encoding  
10 human CTLA-8 (B18). IFN $\gamma$  was used as a positive control angiostatic (i.e., angiogenesis inhibition) activity. These data demonstrate that human CTLA-8 (B18) inhibits angiogenesis.

#### Example 4

##### Hematopoietic Activity of Human CTLA-8

The hematopoietic activity of human CTLA-8 (B18) expressed *in vivo* was examined by construction of a recombinant adenovirus vector.

The B18 cDNA in the expression plasmid Adori 2-12 B18 was driven by the cytomegalovirus(CMV) immediate early promoter and enhancer.

20 The Adori 2-12 vector was created by addition of an SV40 origin and enhancer to a known adenovirus vector (Barr et al., Gene Therapy 1:51 (1994); Davidson et al., Nature Genetics 3:219 (1993)). The HindIII/BamHI fragment encoding the SV40 origin and enhancer was isolated from the pMT2 mammalian expression vector, blunted with Klenow and cloned into the NotI site (blunted with Klenow) of the Ad5  
25 expression vector.

The vector was derived by digesting pNOT-B18 cDNA with *SalI*, filling in the 5' overhang with Klenow to generate a blunt end and digesting with *EcoRI* to isolate the B18 cDNA. The blunted- *EcoRI* B18 fragment was inserted into the restriction sites *EcoRV-EcoRI* of the adenovirus vector Adori 2-12. The CMV-B18 expression  
30 cassette was located downstream of the SV40 origin and enhancer, and 0-1 map units of the left hand end of the adenovirus type 5(Ad5). The SV40 splice donor and

acceptor were located between the CMV promoter and B18 cDNA. Following the insert was SV40 poly A site, 9-16 map units of Ad5 and the puc 19 origin.

A recombinant adenovirus was generated by homologous recombination in 293 cells. *AscI* linearized Adori 2-12 B18 and *ClaI* digested AdCMVlacZ were introduced  
5 into the 293 cells using lipofectamine. Recombinant adenovirus virus was isolated and amplified on 293 cells. The virus was released from infected 293 cells by three cycles of freeze-thawing. The virus was further purified by two cesium chloride centrifugation gradients and dialyzed against PBS 4°C. Following dialysis of the virus glycerol was added to a concentration of 10 % and the virus was stored at - 70 °C until  
10 use. The virus was characterized by expression of the transgene, plaque forming units on 293 cells, particles/ml and Southern analysis of the virus.

A single dose of  $5 \times 10^{10}$  particles of recombinant adenovirus encoding B18 was injected into the tail vein of male C57/bl6 mice, age 7-8 weeks. Control mice received an adenovirus encoding B-galactosidase. Four mice from each experimental  
15 group were killed on day 7 and 14. Blood was collected and automated hematologic analysis was performed using a Baker 9000. Differential counts were performed on blood smears. Tissue was harvested, fixed in formalin, and stained with hematoxylin and eosin for histopathology. In the first set of experiments, serum and tissues were analyzed 7 and 14 days post injection. A slight increase in peripheral platelet counts  
20 were observed. The animals that received B18 exhibited a slight increase in spleen size. Macroscopic analysis of the spleen showed an increase in splenic extramedullary hematopoiesis on day 7 compared to the control. These results showed a hematopoietic growth activity associated with B18.

In a second set of experiments  $5 \times 10^{10}$  particles of recombinant adenovirus encoding B18 were injected into the tail vein of male C57/bl6 mice, age 17-18 weeks.  
25 Control mice received an adenovirus encoding B-galactosidase. Blood samples were collected via retro-orbital sinus on days 2, 5, 7, 10, 14, and 21. The hematologic analyses were performed on the Baker 9000 automated cell counter with murine-specific settings. Analyses included WBC, RBC, HCT and PLT counts. Blood smears  
30 were prepared and stained with Wright-Geimsa for WBC differentials based on a 100 cellcount. Reticulocytes and reticulated platelets were quantitated using flow

cytometry. Four mice from each group were killed on days 7, 14, and 21. In addition to peripheral blood analysis, serum was collected via cardiac puncture for quantitation of systemic IL-6 using a commercial kit (Endogen). Spleen and liver were collected for histopathology, spleen and bone marrow hematopoietic progenitors were quantitated, and bone marrow smears were prepared and stained with Wright-Geimsa for cell counts.

Administration of adenovirus encoding B18 resulted in a marked increase in peripheral blood neutrophils and WBC (Fig. 4). Maximum increases in neutrophils were observed at day 5 and day 7. The control mice showed little difference at day 5 and day 7. Peripheral blood neutrophils were similar in the control mice and mice that received B18 at day 21. In both the B18 and control groups an increase in white blood cells was also observed. The mice that received B18 had a greater increase in WBC between day 2 and day 7. By Day 21 a more pronounced increase was observed in the B-gal group. No other changes in cellular chemistries were observed (Table I).

Bone marrow cellularity was calculated from pooled femurs in each group (Table III). No significant differences were observed in either group. No significant changes were observed in bone marrow hematopoietic progenitors from day 7, 14, and 21. The CFU-GM, BFU-E and CFU-MEG in the B18 mice were similar to the B-gal control (Table II).

Administration of the adenovirus encoding B18 resulted in an increase in CFU-GM (myeloid) and BFU-E (erythroid) progenitors in the spleen compared to animals that received the B-gal virus on day 7. The increase in progenitors in the B18 mice was 11-fold in CFU-GM and a 52-fold in BFU-E (Table II). There was a 2-fold increase in CFU-MEG at day 7 for the B18 mice. By day 21 no significant differences were observed in splenic CFU-MEG or BFU-E between the groups (Table II). A 3-fold decrease in CFU-GM was observed in mice that received adenovirus encoding B18. A slight increase in spleen size at day 7 was observed in the B18 group. This is consistent with an increase in splenic cellularity. By day 14 and day 21 spleen weights were similar to the control group (Table III). Macroscopic analysis of the spleen showed an increase in splenic extramedullary hematopoiesis of the B18 mice on day 7 compared to the control.



The bone marrow myeloid: erythroid ratios (Table IV) suggest a granulocytic hyperplasia with a possible erythroid hypoplasia in mice that received adenovirus B18 on day 7. By day 21 the ratio in the B-gal group was higher. No changes were observed in the IL6 serum levels.

- 5           These results show a hematopoietic activity associated with the administration of adenovirus encoding B18 (human CTLA-8). Increases in neutrophils and white blood cells were observed at day 7 in animals that received B18 adenovirus. The data showed that B18 resulted in increase in splenic CFU-GM and BFU-E 7 days post administration compared to the control animals. Splenic extramedullary  
10 hematopoiesis on day 7 support that B18 exhibits a hematopoietic growth activity. These data suggest that B18 may mobilize early hematopoietic precursors.

Table I: Peripheral hematology for day 2, 5, 10, 14, and 21.

Study AS4-4B...B18 (Platelets) Day 2_4-25-96															
Group A	WBC x10 <sup>3</sup> /uL	Neuts %	ANC x10 <sup>3</sup> /uL	Lymphs %	ALC x10 <sup>3</sup> /uL	Eos %	Monos %	RBC x10 <sup>6</sup> /uL	Retics %	Abs Retics x10 <sup>6</sup> /uL	HCT %	PLT x10 <sup>3</sup> /uL	RPR %	Abs RPR x10 <sup>3</sup> /uL	
B-Gal #1	5.4	40	2.16	54	2.92	0	6	10.88	3.65	0.40	48.0	636	11.94	99.82	
B-Gal #2	7.4	25	1.85	65	4.81	3	7	12.34	2.04	0.25	56.6	900	10.10	90.90	
B-Gal #3	6.8	40	2.72	52	3.54	2	6	11.26	3.26	0.37	51.6	894	9.77	87.34	
B-Gal #4	6.8	23	2.02	64	5.63	1	12	12.00	2.55	0.31	54.8	840	10.63	89.29	
AVG	7.1	32.0	2.19	58.8	4.22	1.5	7.8	11.62	2.88	0.33	52.8	868	10.61	91.84	
SEM	0.7	4.6	0.19	3.4	0.61	0.6	1.4	0.33	0.36	0.03	1.9	17	0.48	2.76	
B18 #1	11.4	59	6.73	31	3.53	1	9	11.16	4.88	0.54	52.4	1242	14.92	185.31	
B18 #2	9.2	30	2.76	62	5.70	3	5	10.14	3.97	0.40	48.0	632	10.90	68.89	
B18 #3	5.0	51	2.55	40	2.00	0	9	11.16	3.23	0.36	51.2	832	11.18	93.02	
B18 #4	6.4	41	2.62	55	3.52	0	4	10.80	3.09	0.33	49.2	904	17.31	156.48	
AVG	8.0	45.3	3.67	47.0	3.69	1.0	6.8	10.82	3.79	0.41	50.2	903	13.58	125.92	
SEM	1.4	6.3	1.02	7.0	0.76	0.7	1.3	0.24	0.41	0.05	1.0	127	1.55	27.07	

Study AS4-4B...B18 (Platelets) Day 5_4-28-96															
Group B	WBC x10 <sup>3</sup> /uL	Neuts %	ANC x10 <sup>3</sup> /uL	Lymphs %	ALC x10 <sup>3</sup> /uL	Eos %	Monos %	RBC x10 <sup>6</sup> /uL	Retics %	Abs Retics x10 <sup>6</sup> /uL	HCT %	PLT x10 <sup>3</sup> /uL	RPR %	Abs RPR x10 <sup>3</sup> /uL	
B-Gal #1	7.6	14	1.06	78	5.93	3	5	11.26	5.25	0.59	52.4	1082	15.51	167.82	
B-Gal #2	10.6	20	2.12	78	8.27	1	1	10.72	4.71	0.50	49.4	994	17.37	172.66	
B-Gal #3	6.8	18	1.51	69	6.07	2	11	11.12	3.40	0.38	51.2	916	9.55	87.48	
B-Gal #4	10.8	38	4.10	58	6.28	0	4	10.22	6.21	0.83	47.0	1092	13.93	152.12	
AVG	9.5	22.5	2.20	70.8	6.63	1.5	5.3	10.83	4.89	0.53	50.0	1021	14.09	145.02	
SEM	0.8	5.3	0.57	4.8	0.55	0.6	2.1	0.23	0.59	0.06	1.2	41	1.67	19.67	
B18 #1	14.8	18	2.66	71	10.51	1	10	12.66	2.31	0.29	57.0	1204	7.57	91.14	
B18 #2	14.2	37	5.25	53	7.53	2	8	9.80	3.32	0.33	44.8	888	14.33	127.25	
B18 #3	12.8	30	3.84	59	7.55	1	10	12.12	4.12	0.50	53.6	1134	10.15	115.10	
B18 #4	16.0	58	9.28	37	5.92	0	5	11.04	3.93	0.43	50.8	1166	15.75	183.65	
AVG	14.5	35.8	6.26	55.0	7.88	1.0	8.3	11.41	3.42	0.39	52.0	1098	11.95	129.28	
SEM	0.7	8.4	1.44	7.1	0.96	0.4	1.2	0.63	0.41	0.05	2.8	71	1.88	19.61	

Study AS4-4B...B18 (Platelets) Day 7_4-30-96															
Group C	WBC x10 <sup>3</sup> /uL	Neuts %	ANC x10 <sup>3</sup> /uL	Lymphs %	ALC x10 <sup>3</sup> /uL	Eos %	Monos %	RBC x10 <sup>6</sup> /uL	Retics %	Abs Retics x10 <sup>6</sup> /uL	HCT %	PLT x10 <sup>3</sup> /uL	RPR %	Abs RPR x10 <sup>3</sup> /uL	
B-Gal #1	15.2	14	2.13	69	10.49	1	16	11.04	3.54	0.39	50.8	862	12.46	107.41	
B-Gal #2	14.0	12	1.68	61	11.34	0	7	11.38	5.05	0.57	52.6	1104	14.91	164.61	
B-Gal #3	14.8	14	2.07	73	10.80	1	12	10.92	5.42	0.59	49.6	952	11.49	109.58	
AVG	14.7	13.3	1.94	74.3	10.88	0.7	11.7	11.11	4.67	0.52	51.0	973	12.95	127.13	
SEM	0.4	0.7	0.14	3.5	0.25	0.3	2.5	0.14	0.58	0.06	0.9	71	1.02	18.75	
B18 #1	18.4	33	6.40	62	12.03	0	5	10.14	2.93	0.30	45.2	864	12.80	110.59	
B18 #2	25.4	39	9.81	53	13.46	0	8	9.46	8.05	0.57	43.6	1288	12.49	160.87	
B18 #3	23.6	44	10.38	50	11.80	0	8	9.74	5.17	0.50	44.4	1076	15.41	165.81	
B18 #4	12.8	15	1.92	75	9.60	0	10	9.54	6.26	0.60	43.4	1136	15.88	180.40	
AVG	20.3	32.8	7.15	60.0	11.72	0.0	7.3	9.72	6.10	0.49	44.2	1091	14.15	154.42	
SEM	2.8	6.3	1.96	5.6	0.80	0.0	1.1	0.15	0.76	0.07	0.4	88	0.87	15.19	

Study AS4-4B...B18 (Platelets) Day 10_5-3-96															
Group A	WBC x10 <sup>3</sup> /uL	Neuts %	ANC x10 <sup>3</sup> /uL	Lymphs %	ALC x10 <sup>3</sup> /uL	Eos %	Monos %	RBC x10 <sup>6</sup> /uL	Retics %	Abs Retics x10 <sup>6</sup> /uL	HCT %	PLT x10 <sup>3</sup> /uL	RPR %	Abs RPR x10 <sup>3</sup> /uL	
B-Gal #1	18.8	17	3.16	69	12.83	3	11	10.22	12.41	1.27	46.8	1460	16.20	236.52	
B-Gal #2	13.2	16	2.11	79	10.43	1	4	10.48	8.00	0.63	48.8	1128	14.48	163.33	
B-Gal #3	19.6	16	3.14	74	14.50	0	10	10.72	6.25	0.67	49.4	1338	16.58	221.84	
B-Gal #4	18.8	21	3.91	72	13.39	3	4	10.44	7.59	0.79	48.4	1068	14.35	153.26	
AVG	17.5	17.5	3.08	73.5	12.79	1.8	7.3	10.47	8.06	0.84	48.4	1249	15.40	193.74	
SEM	1.5	1.2	0.37	2.1	0.86	0.5	1.9	0.10	1.49	0.15	0.6	91	0.58	20.78	
B18 #1	14.2	33	4.69	56	7.95	5	6	8.70	11.97	1.04	39.2	1760	14.49	255.02	
B18 #2	17.6	35	6.16	57	10.03	1	7	9.04	9.46	0.86	42.0	1104	18.88	208.44	
B18 #3	18.2	39	6.32	57	9.23	1	3	4.74	16.77	0.79	22.4	894	29.19	260.96	
B18 #4	14.2	25	3.55	66	9.37	1	8	9.30	9.93	0.92	42.0	1416	16.81	238.03	
AVG	16.6	33.0	6.18	59.0	9.15	2.0	6.0	7.95	12.04	0.90	36.4	1294	19.84	240.61	
SEM	0.8	2.9	0.66	2.3	0.43	1.0	1.1	1.08	1.57	0.05	4.7	189	3.24	11.77	

Study AS4-4B...B18 (Platelets) Day 14_5-7-96															
Group B	WBC x10 <sup>3</sup> /uL	Neuts %	ANC x10 <sup>3</sup> /uL	Lymphs %	ALC x10 <sup>3</sup> /uL	Eos %	Monos %	RBC x10 <sup>6</sup> /uL	Retics %	Abs Retics x10 <sup>6</sup> /uL	HCT %	PLT x10 <sup>3</sup> /uL	RPR %	Abs RPR x10 <sup>3</sup> /uL	
B-Gal #1	17.8	18	3.20	74	13.17	0	8	10.88	5.97	0.65	50.8	1360	11.03	150.01	
B-Gal #2	20.4	26	5.30	66	13.48	1	7	10.92	7.07	0.77	50.8	1616	8.16	132.19	
B-Gal #3	16.0	7	1.12	90	14.40	1	3	11.36	6.41	0.73	52.8	1298	7.36	95.53	
B-Gal #4	18.0	36	6.48	57	10.26	1	6	9.30	7.82	0.71	43.0	1672	10.05	168.04	
AVG	18.1	21.8	4.03	71.8	12.82	0.8	6.0	10.61	6.77	0.71	49.4	1487	9.16	138.44	
SEM	0.9	6.1	1.18	7.0	0.89	0.3	1.1	0.45	0.36	0.03	2.2	93	0.84	15.48	
B18 #1	15.4	9	1.39	81	12.47	1	9	10.62	5.74	0.61	46.2	1262	9.51	120.02	
B18 #2	15.4	31	4.77	58	8.93	2	9	9.78	10.33	1.01	44.6	1092	14.29	156.05	
B18 #3	13.4	42	5.63	39	5.23	0	19	10.34	4.99	0.52	48.8	1378	15.79	217.27	
B18 #4	11.6	57	6.61	34	3.94	2	7	9.38	5.57	0.52	43.0	1092	16.66	181.93	
AVG	14.0	34.8	4.60	53.0	7.64	1.3	11.0	10.03	6.66	0.66	45.6	1206	14.06	168.82	
SEM	0.9	10.1	1.14	10.7	1.93	0.5	2.7	0.26	1.23	0.12	1.1	70	1.59	20.54	

Study AS4-4B...B18 (Platelets) Day 21_5-14-96															
Group A	WBC x10 <sup>3</sup> /uL	Neuts %	ANC x10 <sup>3</sup> /uL	Lymphs %	ALC x10 <sup>3</sup> /uL	Eos %	Monos %	RBC x10 <sup>6</sup> /uL	Retics %	Abs Retics x10 <sup>6</sup> /uL	HCT %	PLT x10 <sup>3</sup> /uL	RPR %	Abs RPR x10 <sup>3</sup> /uL	
B-Gal #1	25.4	23	5.84	67	17.02	0	10	9.22	6.15	0.75	42.8	1776	9.61	170.87	
B-Gal #2	19.8	19	3.72	69	13.32	0	12	9.50	9.95	0.95	44.4	1662	9.44	156.89	
B-Gal #3	27.8	11	3.04	82	22.63	3	4	9.74	8.84	0.86	45.8	1684	11.45	192.82	
B-Gal #4	28.0	13	3.64	83	23.24	0	4	9.04	7.54	0.68	41.6	1346	10.48	141.06	
AVG	25.2	16.5	4.06	75.3	19.10	0.8	7.5	9.38	8.62	0.81	43.7	1617	10.25	185.36	
SEM	1.9	2.8	0.61	4.2	2.33	0.8	2.1	0.15	0.52	0.08	0.9	94	0.46	10.97	
B18 #1	18.8	9	1.67	83	15.44	1	7	9.84	7.40	0.73	43.8	1642	8.37	137.44	
B18 #2	16.2	48	7.78	45	7.29	1	6	11.10	7.97	0.88	51.4	1970	10.78	212.37	
B18 #3	14.8	53	7.84	42	6.22	0	5	7.52	20.26	1.52	38.0	1568	10.50	164.64	
B18 #4	15.2	18	2.74	74	11.25	2	6	9.84	7.02	0.68	43.0	1422	7.60	108.07	
AVG	16.2	32.0	5.01	61.0	10.05	1.0	6.0	9.53	10.66	0.95	44.1	1651	9.31	155.63	
SEM	0.9	10.9	1.63	10.3	2.10	0.4	0.4	0.74	3.21	0.20	2.8	116	0.78	22.16	

Table II: Bone marrow and Splenic Hematopoietic Progenitors

	CFU-MEG		CFU-GM		BFU-E	
	B-Gal	B18	B-Gal	B18	B-Gal	B18
Bone Marrow*						
Day 7	16.0 ± 3.5	15.7 ± 3.1	307 ± 117	241 ± 78	51 ± 19	25 ± 11
Day 14	10.7 ± 2.3	15.3 ± 1.2	233 ± 15	373 ± 35	30 ± 10	60 ± 30
Day 21	5.7 ± 0.6	6.7 ± 3.1	170 ± 17	160 ± 27	40 ± 10	27 ± 6
Spleen**						
Day 7	9.3 ± 1.6	19.5 ± 1.5	27 ± 3	298 ± 6	1.3 ± 1.2	68 ± 10
Day 14	9.7 ± 0.6	12.7 ± 0.6	267 ± 32	197 ± 21	33 ± 6	10 ± 10
Day 21	17.0 ± 1.0	19.3 ± 2.5	187 ± 6	73 ± 15	23 ± 6	23 ± 6

Hematopoietic precursors were determined from pooled spleen and bone marrow samples from four animals in each group. For quantitation of CFU-GM and BFU-E, either  $1 \times 10^4$  bone marrow cells or  $1 \times 10^5$  spleen cells were added to complete alpha methylcellulose medium (0.9% methylcellulose in alpha medium, 30% fetal bovine serum, 1% bovine serum albumin, 10-4M 2-mercaptoethanol, 2 mM L-glutamine, 2% murine spleen cell conditioned medium, and 3 U/mL erythropoietin) and aliquoted into 35 mm tissue culture dishes in a final volume of 1.0 mL. Cultures were incubated for 7 days at 37°C, and 5% CO<sub>2</sub>. Microscopic colonies were defined as clusters of 50 or more cells. For quantitation of CFU-MEG, either  $1 \times 10^5$  bone marrow cells or  $1 \times 10^6$  spleen cells were added to complete alpha methylcellulose medium and incubated as described above. Megakaryocyte colonies were defined as a group of 3 or more cells.

\*Bone marrow progenitors are represented as mean ± sd number of colonies per  $10^5$  cells.

\*\*Spleen progenitors are represented as mean ± sd number of colonies per  $10^6$  cells.

Table III: Spleen Weights and Femur Cellularity

Spleen Wt. (Mg)	B-Gal	B18		Femur Cellularity (x10 <sup>6</sup> )	B-Gal	B18
Day 7	187 ± 19	224 ± 29		Day 7	28	23
Day 14	175 ± 13	170 ± 10		Day 14	28	27
Day 21	174 ± 21	151 ± 27		Day 21	28	26

Spleen weights were determined at time of sacrifice are represented as means ± sd from four animals.

Table IV: Bone Marrow Myeloid:Erythroid Ratios

Group	Mouse #	Day 7	Day 14	Day 21
B-gal	1	1.43	2.12	5.78
	2	0.91	2.46	5.83
	3	1.62	1.03	3.66
	4		5.44	4.82
<b>AVG</b>		<b>1.32</b>	<b>2.76</b>	<b>5.02</b>
<b>SD</b>		<b>0.37</b>	<b>0.37</b>	<b>1.89</b>
B-gal	1	5.59	2.01	2.02
	2	6.51	1.25	2.13
	3	5.49	1.58	1.81
	4	0.50	2.51	2.92
<b>AVG</b>		<b>4.52</b>	<b>1.86</b>	<b>2.22</b>
<b>SD</b>		<b>1.29</b>	<b>2.72</b>	<b>0.56</b>

All entries represent the number of myeloid cells per 1 erythroid cell. Normal mouse ratios are approximately 1:1 to 2:1.

#### Example 5

#### Additional Experiments Relating to Hematopoietic Activity of Human CTLA-8

B18 (human CTLA-8) was tested for the ability to induce production of factors having hematopoietic activity in a factor-dependent cell proliferation assay using the

human erythroleukemic cell line, TF-1 (Kitamura et al., J. Cell Physiol. 140:323 (1989)). The cells were initially grown in the presence of rhGMCSF (100 U/ml). The cells were fed three days prior to setting up the assay. The assay conditions were as follows:

5	cells/well	5000/200 $\mu$ l
	incubation time	3 days
	pulse time	4 hours
	amount of tritiated thymidine	0.5 $\mu$ Ci/well
	counting time	1 minute
10	replicates	2

B18 alone, conditioned medium (CM) from B18 induced HS-5 cells were assayed. Buffer alone, CM from HS-5 cells induced with buffer and CM from uninduced HS-5 cells were assayed as controls. Results are shown in Fig. 5. B18 (human CTLA-8) demonstrated an ability to induce production of factors which induced TF-1 proliferation. This activity was substantially eliminated by the addition of anti-GMCSF antibodies. These data demonstrate that human CTLA-8 (B18) is able to induce hematopoiesis. Particularly, without being bound by any theory, it appears that human CTLA-8 (B18) induces production of GM-CSF and/or IL-3.

20

#### Example 6

##### Ability of Human CTLA-8 to Induce Production of IL-6 and IL-8

MRC5 cells were incubated in the presence of human CTLA-8 (B18) and production of IL-6 and IL-8 were measured. Herpes CTLA-8 (IL-17) was used as a positive control. Applicants' human CTLA-8 (B18) demonstrated titratable production of both IL-6 and IL-8 (see Figs. 6 and 7).

25

All patent and literature references cited herein are incorporated by reference as if fully set forth.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: Jacobs, Kenneth  
Kelleher, Kerry  
Carlin, McKeough  
Goldman, Samuel  
Pittman, Debra  
Mi, Sha  
Neben, Steven  
Giannotti, JoAnn  
Golden Fleet, Margaret
- (ii) TITLE OF INVENTION: Human CTLA-8 and Uses of CTLA-8-Related Proteins
- (iii) NUMBER OF SEQUENCES: 9
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Genetics Institute, Inc.
  - (B) STREET: 87 CambridgePark Drive
  - (C) CITY: Cambridge
  - (D) STATE: Massachusetts
  - (E) COUNTRY: USA
  - (F) ZIP: 02140
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER:
  - (B) FILING DATE:
  - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: Brown, Scott A.
  - (B) REGISTRATION NUMBER: 32,724
  - (C) REFERENCE/DOCKET NUMBER: GI5262
- (ix) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: (617) 498-8224
  - (B) TELEFAX: (617) 876-5851

## (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 813 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (ix) FEATURE:
  - (A) NAME/KEY: CDS
  - (B) LOCATION: 56..544

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```

GGGAAGATAC ATTACAGAA AGAGCTTCCT GCACAAAGTA AGCCACCAGC GCAAC ATG      58
                                   Met
                                   1

ACA GTG AAG ACC CTG CAT GGC CCA GCC ATG GTC AAG TAC TTG CTG CTG      106
Thr Val Lys Thr Leu His Gly Pro Ala Met Val Lys Tyr Leu Leu
      5                               10                               15

TCG ATA TTG GGG CTT GCC TTT CTG AGT GAG GCG GCA GCT CGG AAA ATC      154
Ser Ile Leu Gly Leu Ala Phe Leu Ser Glu Ala Ala Arg Lys Ile
      20                               25                               30

CCC AAA GTA GGA CAT ACT TTT TTC CAA AAG CCT GAG AGT TGC CCG CCT      202
Pro Lys Val Gly His Thr Phe Gln Lys Pro Glu Ser Cys Pro Pro
      35                               40                               45

GTG CCA GGA GGT AGT ATG AAG CTT GAC ATT GGC ATC ATC AAT GAA AAC      250
Val Pro Gly Gly Ser Met Lys Leu Asp Ile Gly Ile Ile Asn Glu Asn
      50                               55                               60                               65

CAG CGC GTT TCC ATG TCA CGT AAC ATC GAG AGC CGC TCC ACC TCC CCC      298
Gln Arg Val Ser Met Ser Arg Asn Ile Glu Ser Arg Ser Thr Ser Pro
      70                               75                               80

TGG AAT TAC ACT GTC ACT TGG GAC CCC AAC CGG TAC CCC TCG GAA GTT      346
Trp Asn Tyr Thr Val Thr Trp Asp Pro Asn Arg Tyr Pro Ser Glu Val
      85                               90                               95

GTA CAG GCC CAG TGT AGG AAC TTG GGC TGC ATC AAT GCT CAA GGA AAG      394
Val Gln Ala Gln Cys Arg Asn Leu Gly Cys Ile Asn Ala Gln Gly Lys
      100                              105                              110

GAA GAC ATC TCC ATG AAT TCC GTT CCC ATC CAG CAA GAG ACC CTG GTC      442
Glu Asp Ile Ser Met Asn Ser Val Pro Ile Gln Gln Glu Thr Leu Val
      115                              120                              125

GTC CGG AGG AAG CAC CAA GGC TGC TCT GTT TCT TTC CAG TTG GAG AAG      490
Val Arg Arg Lys His Gln Gly Cys Ser Val Ser Phe Gln Leu Glu Lys
      130                              135                              140                              145

GTG CTG GTG ACT GTT GGC TGC ACC TGC GTC ACC CCT GTC ATC CAC CAT      538
Val Leu Val Thr Val Gly Cys Thr Cys Val Thr Pro Val Ile His His
      150                              155                              160

GTG CAG TAAGAGGTGC ATATCCACTC AGCTGAAGAA GCTGTAGAAA TGCCACTCCT      594
Val Gln

TACCCAGTGC TCTGCAACAA GTCCTGTCTG ACCCCCAATT CCCTCCACTT CACAGGACTC      654

TTAATAAGAC CTGCACGGAT GGAAACAGAA AATATTCACA ATGTATGTGT GTATGTACTA      714

CACTTTATAT TTGATATCTA AAATGTTAGG AGAAAAATTA ATATATTCAG TGCTAATATA      774

ATAAAGTATT AATAATTTAA AAATAAAAAA AAAAAAAA      813

```

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 163 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Thr Val Lys Thr Leu His Gly Pro Ala Met Val Lys Tyr Leu Leu  
 1 5 10 15  
 Leu Ser Ile Leu Gly Leu Ala Phe Leu Ser Glu Ala Ala Ala Arg Lys  
 20 25 30  
 Ile Pro Lys Val Gly His Thr Phe Phe Gln Lys Pro Glu Ser Cys Pro  
 35 40 45  
 Pro Val Pro Gly Gly Ser Met Lys Leu Asp Ile Gly Ile Ile Asn Glu  
 50 55 60  
 Asn Gln Arg Val Ser Met Ser Arg Asn Ile Glu Ser Arg Ser Thr Ser  
 65 70 75 80  
 Pro Trp Asn Tyr Thr Val Thr Trp Asp Pro Asn Arg Tyr Pro Ser Glu  
 85 90 95  
 Val Val Gln Ala Gln Cys Arg Asn Leu Gly Cys Ile Asn Ala Gln Gly  
 100 105 110  
 Lys Glu Asp Ile Ser Met Asn Ser Val Pro Ile Gln Gln Glu Thr Leu  
 115 120 125  
 Val Val Arg Arg Lys His Gln Gly Cys Ser Val Ser Phe Gln Leu Glu  
 130 135 140  
 Lys Val Leu Val Thr Val Gly Cys Thr Cys Val Thr Pro Val Ile His  
 145 150 155 160  
 His Val Gln

## (2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 461 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

- (ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 6..455

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CCACC ATG TGC CTG ATG CTG TTG CTG CTA CTG AAC CTG GAG GCT ACA 47  
 Met Cys Leu Met Leu Leu Leu Leu Asn Leu Glu Ala Thr  
 1 5 10  
 GTG AAG GCA GCG GTA CTC ATC CCT CAA AGT TCA GTG TGT CCA AAC GCC 95  
 Val Lys Ala Ala Val Leu Ile Pro Gln Ser Ser Val Cys Pro Asn Ala  
 15 20 25 30  
 GAG GCC AAT AAC TTT CTC CAG AAC GTG AAG GTC AAC CTG AAA GTC ATC 143  
 Glu Ala Asn Asn Phe Leu Gln Asn Val Lys Val Asn Leu Lys Val Ile  
 35 40 45



AAC TCC CTT AGC TCA AAA GCG AGC TCG AGA AGG CCC TCA GAT TAC CTC	191
Asn Ser Leu Ser Ser Lys Ala Ser Ser Arg Arg Pro Ser Asp Tyr Leu	
50 55 60	
AAC CGT TCC ACT TCA CCC TGG ACT CTG AGC CGC AAT GAG GAC CCT GAT	239
Asn Arg Ser Thr Ser Pro Trp Thr Leu Ser Arg Asn Glu Asp Pro Asp	
65 70 75	
AGA TAT CCT TCT GTG ATC TGG GAG GCA CAG TGC CGC CAC CAG CGC TGT	287
Arg Tyr Pro Ser Val Ile Trp Glu Ala Gln Cys Arg His Gln Arg Cys	
80 85 90	
GTC AAC GCT GAG GGG AAG TTG GAC CAC CAC ATG AAT TCT GTT CTC ATC	335
Val Asn Ala Glu Gly Lys Leu Asp His His Met Asn Ser Val Leu Ile	
95 100 105 110	
CAG CAA GAG ATA CTA GTC CTG AAG AGG GAG CCT GAG AAG TGC CCC TTC	383
Gln Gln Glu Ile Leu Val Leu Lys Arg Glu Pro Glu Lys Cys Pro Phe	
115 120 125	
ACT TTC CGG GTG GAG AAG ATG CTG GTG GGC GTG GGC TGC ACC TGC GTT	431
Thr Phe Arg Val Glu Lys Met Leu Val Gly Val Gly Cys Thr Cys Val	
130 135 140	
TCC TCT ATT GTC CGC CAT GCG TCC TAATAA	461
Ser Ser Ile Val Arg His Ala Ser	
145 150	

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 150 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Cys Leu Met Leu Leu Leu Leu Leu Asn Leu Glu Ala Thr Val Lys	
1 5 10 15	
Ala Ala Val Leu Ile Pro Gln Ser Ser Val Cys Pro Asn Ala Glu Ala	
20 25 30	
Asn Asn Phe Leu Gln Asn Val Lys Val Asn Leu Lys Val Ile Asn Ser	
35 40 45	
Leu Ser Ser Lys Ala Ser Ser Arg Arg Pro Ser Asp Tyr Leu Asn Arg	
50 55 60	
Ser Thr Ser Pro Trp Thr Leu Ser Arg Asn Glu Asp Pro Asp Arg Tyr	
65 70 75 80	
Pro Ser Val Ile Trp Glu Ala Gln Cys Arg His Gln Arg Cys Val Asn	
85 90 95	
Ala Glu Gly Lys Leu Asp His His Met Asn Ser Val Leu Ile Gln Gln	
100 105 110	
Glu Ile Leu Val Leu Lys Arg Glu Pro Glu Lys Cys Pro Phe Thr Phe	
115 120 125	
Arg Val Glu Lys Met Leu Val Gly Val Gly Cys Thr Cys Val Ser Ser	
130 135 140	

Ile Val Arg His Ala Ser  
145 150

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 459 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(ix) FEATURE:

- (A) NAME/KEY: CDS  
 (B) LOCATION: 1..453

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

ATG ACA TTT AGA ATG ACT TCA CTT GTG TTA CTT CTG CTG CTG AGC ATA	48
Met Thr Phe Arg Met Thr Ser Leu Val Leu Leu Leu Leu Ser Ile	
1 5 10 15	
GAT TGT ATA GTA AAG TCA GAA ATA ACT AGT GCA CAA ACC CCA AGA TGC	96
Asp Cys Ile Val Lys Ser Glu Ile Thr Ser Ala Gln Thr Pro Arg Cys	
20 25 30	
TTA GCT GCT AAC AAT AGC TTT CCA CGG TCT GTG ATG GTT ACT TTG AGC	144
Leu Ala Ala Asn Asn Ser Phe Pro Arg Ser Val Met Val Thr Leu Ser	
35 40 45	
ATC CGT AAC TGG AAT ACC AGT TCT AAA AGG GCT TCA GAC TAC TAC AAT	192
Ile Arg Asn Trp Asn Thr Ser Ser Lys Arg Ala Ser Asp Tyr Tyr Asn	
50 55 60	
AGA TCT ACG TCT CCT TGG ACT CTC CAT CGC AAT GAA GAT CAA GAT AGA	240
Arg Ser Thr Ser Pro Trp Thr Leu His Arg Asn Glu Asp Gln Asp Arg	
65 70 75 80	
TAT CCC TCT GTG ATT TGG GAA GCA AAG TGT CGC TAC TTA GGA TGT GTT	288
Tyr Pro Ser Val Ile Trp Glu Ala Lys Cys Arg Tyr Leu Gly Cys Val	
85 90 95	
AAT GCT GAT GGG AAT GTA GAC TAC CAC ATG AAC TCA GTC CCT ATC CAA	336
Asn Ala Asp Gly Asn Val Asp Tyr His Met Asn Ser Val Pro Ile Gln	
100 105 110	
CAA GAG ATT CTA GTG GTG CGC AAA GGG CAT CAA CCC TGC CCT AAT TCA	384
Gln Glu Ile Leu Val Val Arg Lys Gly His Gln Pro Cys Pro Asn Ser	
115 120 125	
TTT AGG CTA GAG AAG ATG CTA GTG ACT GTA GGC TGC ACA TGC GTT ACT	432
Phe Arg Leu Glu Lys Met Leu Val Thr Val Gly Cys Thr Cys Val Thr	
130 135 140	
CCC ATT GTT CAC AAT GTA GAC TAAAAG	459
Pro Ile Val His Asn Val Asp	
145 150	

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 151 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

```

Met Thr Phe Arg Met Thr Ser Leu Val Leu Leu Leu Leu Ser Ile
 1           5           10           15
Asp Cys Ile Val Lys Ser Glu Ile Thr Ser Ala Gln Thr Pro Arg Cys
          20           25           30
Leu Ala Ala Asn Asn Ser Phe Pro Arg Ser Val Met Val Thr Leu Ser
          35           40           45
Ile Arg Asn Trp Asn Thr Ser Ser Lys Arg Ala Ser Asp Tyr Tyr Asn
          50           55           60
Arg Ser Thr Ser Pro Trp Thr Leu His Arg Asn Glu Asp Gln Asp Arg
          65           70           75           80
Tyr Pro Ser Val Ile Trp Glu Ala Lys Cys Arg Tyr Leu Gly Cys Val
          85           90           95
Asn Ala Asp Gly Asn Val Asp Tyr His Met Asn Ser Val Pro Ile Gln
          100          105          110
Gln Glu Ile Leu Val Val Arg Lys Gly His Gln Pro Cys Pro Asn Ser
          115          120          125
Phe Arg Leu Glu Lys Met Leu Val Thr Val Gly Cys Thr Cys Val Thr
          130          135          140
Pro Ile Val His Asn Val Asp
          145          150

```

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 28 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CACAGGCATA CACAGGAAGA TACATTCA

28

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 25 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

TCTTGCTGGA TGGGAACGGA ATTCA

25

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 28 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ATACATTCAC AGAAGAGCTT CCTGCACA

28

5 What is claimed is:

1. An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

(a) the nucleotide sequence of SEQ ID NO:1 from nucleotide 146 to nucleotide 544;

10 (b) a nucleotide sequence capable of hybridizing to a nucleic acid sequence specified in (a);

(c) a nucleotide sequence varying from the sequence of the nucleotide sequence specified in (a) as a result of degeneracy of the genetic code; and

(d) an allelic variant of the nucleotide sequence specified in (a).

15

2. The polynucleotide of claim 1 wherein said nucleotide sequence encodes a protein having CTLA-8 activity.

3. The polynucleotide of claim 1 wherein said nucleotide sequence is operably linked to an expression control sequence.

20

4. The polynucleotide of claim 1 comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 55 to nucleotide 544.

25 5. A host cell transformed with the polynucleotide of claim 3.

6. The host cell of claim 5, wherein said cell is a mammalian cell.

7. A process for producing a human CTLA-8 protein, said process comprising:

30 (a) growing a culture of the host cell of claim 5 in a suitable culture medium;  
and

(b) purifying the human CTLA-8 protein from the culture.

- 5      8.      An isolated human CTLA-8 protein comprising an amino acid sequence selected from the group consisting of:
- (a)      the amino acid sequence of SEQ ID NO:2;
  - (b)      the amino acid sequence of SEQ ID NO:2 from amino acids 11 to 163;
  - (c)      the amino acid sequence of SEQ ID NO:2 from amino acids 29 to 163;
  - 10      (d)      the amino acid sequence of SEQ ID NO:2 from amino acids 31 to 163; and
  - (e)      fragments of (a), (b), (c) or (d) having CTLA-8 activity.
9.      The protein of claim 8 comprising the amino acid sequence of SEQ ID NO:2.
- 15      10.      The protein of claim 8 comprising the sequence from amino acid 29 to 163 of SEQ ID NO:2.
11.      A pharmaceutical composition comprising a human CTLA-8 protein of claim 8 and a pharmaceutically acceptable carrier.
- 20      12.      A human CTLA-8 protein produced according to the process of claim 7.
13.      A composition comprising an antibody which specifically reacts with a human CTLA-8 protein of claim 8.
- 25      14.      A method of treating a mammalian subject comprising administering a therapeutically effective amount of a composition of claim 11.
15.      A method of treating a mammalian subject comprising administering a therapeutically effective amount of a composition comprising a pharmaceutically acceptable carrier and a protein comprising an amino acid sequence selected from the group consisting of:
- 30      (a)      the amino acid sequence of SEQ ID NO:4;

- 5           (b)     the amino acid sequence of SEQ ID NO:4 from amino acids 18 to 150; and  
          (c)     fragments of (a) or (b) having CTLA-8 activity.

16.     The method of claim 15 wherein said protein comprises the amino acid sequence of SEQ ID NO:4.

10

17.     The method of claim 15 wherein said protein comprises the amino acid sequence of SEQ ID NO:4 from amino acids 18 to 150.

15

18.     A method of treating a mammalian subject administering a therapeutically effective amount of a composition comprising a pharmaceutically acceptable carrier and a protein comprising an amino acid sequence selected from the group consisting of:

- (a)     the amino acid sequence of SEQ ID NO:6;
- (b)     the amino acid sequence of SEQ ID NO:6 from amino acids 19 to 151; and
- (c)     fragments of (a) or (b) having CTLA-8 activity.

20

19.     The method of claim 18 wherein said protein comprises the amino acid sequence of SEQ ID NO:6.

25

20.     The method of claim 18 wherein said protein comprises the amino acid sequence of SEQ ID NO:6 from amino acids 19 to 151.

21.     The polynucleotide of claim 1 comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 86 to nucleotide 544.

30

22.     The protein of claim 8 comprising the sequence from amino acid 11 to 163 of SEQ ID NO:2.

5 23. A method of treating a mammalian subject comprising administering a therapeutically effective amount of a composition comprising a pharmaceutically acceptable carrier and IL-17 or an active fragment thereof.

10 24. The method of claim 14, 15, 18 or 23 wherein said subject is treated to produce an effect selected from the group consisting of inhibition of angiogenesis, inhibition of growth or proliferation of vascular endothelial cells, inhibition of tumor growth, inhibition of angiogenesis-dependent tissue growth, proliferation of myeloid cells or progenitors, proliferation of erythroid cells or progenitors, proliferation of lymphoid cells or progenitors, induction of IFN $\gamma$  production, induction of IL-3 production and induction of GM-CSF  
15 production.

25. The composition of claim 3 wherein said polynucleotide is contained in a vector suitable for *in vivo* expression in a mammalian subject.

20 26. The polynucleotide of claim 1 comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 139 to nucleotide 544.

27. The protein of claim 8 comprising the sequence from amino acid 31 to 163 of SEQ ID NO:2.





2/7




FIGURE 2

3/7




Fig. 3

4/7




Fig. 4

## TF-1 activity in CM from B18 induced HS-5 cells




Fig. 5

IL6 Induction from B18 on MRC5 Cells - 24 & 48hrs




Fig. 6

IL8 Production from B18 on MRC5 Cells - 24 & 48hrs




Fig. 7

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification<sup>6</sup> :</b> <b>C12N 15/19, 15/12, 15/24, C07K 14/52,</b> <b>14/725, 14/54, A61K 38/17, 38/19, 38/20,</b> <b>C12N 5/10, C07K 16/24, 16/28, A61K 48/00</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 97/04097</b> <b>(43) International Publication Date:</b> 6 February 1997 (06.02.97)
<b>(21) International Application Number:</b> PCT/US96/11889 <b>(22) International Filing Date:</b> 18 July 1996 (18.07.96)  <b>(30) Priority Data:</b> 08/504,032 19 July 1995 (19.07.95) US 08/514,014 11 August 1995 (11.08.95) US  <b>(71) Applicant:</b> GENETICS INSTITUTE, INC. [US/US]; 87 Cambridge Park Drive, Cambridge, MA 02140 (US).  <b>(72) Inventors:</b> JACOBS, Kenneth; 151 Beaumont Avenue, Newton, MA 02160 (US). KELLEHER, Kerry; 50 Hurley Circle, Marlborough, MA 01752 (US). CARLIN, McKeough; 16 Chauncy Street #2, Cambridge, MA 02138 (US). GOLDMAN, Samuel; 9 Mohawk Drive, Acton, MA 01720 (US). PITTMAN, Debra; 20 N. Shore Road, Windham, NH 03087 (US). MI, Sha; 4 Vernon Road, Belmont, MA 02178 (US). NEBEN, Steven; 13 Duggan Road, Acton, MA 01720 (US). GIANNOTTI, JoAnn; 409 Arlington Street, Acton, MA 01720 (US). GOLDEN-FLEET, Margaret; 19 Usher Road, Medford, MA 02155 (US).  <b>(74) Agent:</b> BROWN, Scott, A.; Genetics Institute, Inc., 87 CambridgePark Drive, Cambridge, MA 02140 (US).	<b>(81) Designated States:</b> AU, CA, JP, MX, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>  <b>(88) Date of publication of the international search report:</b> 12 September 1997 (12.09.97)	
<b>(54) Title:</b> HUMAN CTLA-8 AND USES OF CTLA-8-RELATED PROTEINS  <b>(57) Abstract</b>  Polynucleotides encoding human CTLA-8 and related proteins are disclosed. Human CTLA-8 proteins and methods for their production are also disclosed. Methods of treatment using human CTLA-8 proteins, rat CTLA-8 proteins and herpesvirus herpes CTLA-8 proteins are also provided.		



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LJ	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 96/11889

A. CLASSIFICATION OF SUBJECT MATTER					
IPC 6	C12N15/19	C12N15/12	C12N15/24	C07K14/52	C07K14/725
	C07K14/54	A61K38/17	A61K38/19	A61K38/20	C12N5/10
	C07K16/24	C07K16/28	A61K48/00		
According to International Patent Classification (IPC) or to both national classification and IPC					
B. FIELDS SEARCHED					
Minimum documentation searched (classification system followed by classification symbols)					
IPC 6	C12N	C07K	A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched					
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)					
C. DOCUMENTS CONSIDERED TO BE RELEVANT					
Category *	Citation of document, with indication, where appropriate, of the relevant passages				Relevant to claim No.
X	WO 95 18826 A (SCHERING CORP ; INST NAT SANTE RECH MED (FR)) 13 July 1995 cited in the application				1-3,5-7, 12, 15-20, 23-25
A	see page 3, line 21 - page 4, line 11 see page 12, line 1 - page 16, line 6 see page 30, line 1-13 Seq.ID:2 see page 53 Seq.ID:4 see page 55 Seq.ID:5 see page 56 Seq.ID:7/8 see page 58 - page 59 --- -/--				11,13,14
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.					
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "A" document member of the same patent family					
Date of the actual completion of the international search			Date of mailing of the international search report		
21 July 1997			25.07.97		
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl, Fax (+ 31-70) 340-3016			Authorized officer  Macchia, G		

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 96/11889

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>THE JOURNAL OF IMMUNOLOGY, vol. 150, no. 12, 15 June 1993, pages 5445-5456, XP002035505 ROUVIER E. ET AL.: "CTLA-8, cloned from an activated T cell, bearing AU-rich messenger RNA instability sequences, and homologous to a Herpesvirus Saimiri gene" cited in the application see page 5445 - page 5446, line 11 see page 5449, right-hand column, line 26 - page 5450 see page 5451; figure 3 see page 5453, left-hand column, line 10 - right-hand column, line 16 ---</p>	15-20
X	<p>JOURNAL OF VIROLOGY, vol. 66, no. 8, August 1992, pages 5047-5058, XP000615399 ALBRECHT J -C ET AL: "PRIMARY STRUCTURE OF THE HERPESVIRUS SAIMIRI GENOME" Seq.ID:1 from nt.552 to nt.217 (reverse orientation) is 61.4% homologous to X64346 from nt.26931 to nt.27266. see page 5048, right-hand column, line 15-18 ORF13 see page 5049; table 1 ---</p>	1
A	<p>WO 97 07198 A (GENETICS INST) 27 February 1997 see page 8, line 15-31 see page 9, line 26 - page 11, line 7 see page 13, line 16-27 see page 18, line 10 - page 19, line 25 see page 25, line 11-15 see page 27, line 5-17 see page 31, line 27-34 Seq.ID:11-12 see page 49 - page 51 ---</p>	18-20
E	<p>WO 97 07198 A (GENETICS INST) 27 February 1997 see page 8, line 15-31 see page 9, line 26 - page 11, line 7 see page 13, line 16-27 see page 18, line 10 - page 19, line 25 see page 25, line 11-15 see page 27, line 5-17 see page 31, line 27-34 Seq.ID:11-12 see page 49 - page 51 ---</p>	1-14,21, 22,24-27
P,X	<p>JOURNAL OF IMMUNOLOGY, vol. 155, no. 12, 15 December 1995, pages 5483-5486, XP000602481 YAO Z ET AL: "HUMAN IL-17: A NOVEL CYTOKINE DERIVED FROM T CELLS" see page 5483 see page 5484; figure 1 ---</p>	1-3,5-7, 23
	<p>---</p>	

-/--

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 96/11889

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	THE JOURNAL OF EXPERIMENTAL MEDICINE, vol. 183, no. 6, 1 June 1996, pages 2593-2603, XP002035506 FOSSIEZ F. ET AL.: "T cell interleukin-17 induces stromal cells to produce proinflammatory and hematopoietic cytokines" see abstract see page 2594; figure 1 ---	1-3,5-7, 15-20, 23,24
P,X	IMMUNITY, vol. 3, no. 6, 1 December 1995, pages 811-821, XP000579309 YAO Z ET AL: "HERPESVIRUS SAIMIRI ENCODES A NEW CYTOKINE, IL-17, WHICH BINDS TO A NOVEL CYTOKINE RECEPTOR" see page 811 see page 815, left-hand column, line 50 - page 818, left-hand column ---	15-20, 23,24
E	WO 96 29408 A (IMMUNEX CORP) 26 September 1996 see page 1, line 1 - page 2, line 33 Seq.ID:8 see page 36 ---	18-20
P,X	GENE, vol. 168, no. 2, 12 February 1996, pages 223-225, XP002035631 YAO ET AL.: "Complete nucleotide sequence of the mouse CTLA8 gene" see page 225; figure 2 -----	1,2

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 96/11889

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 14-20,23,24  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 14-20,2,24 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see continuation-sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 96/ 11889

FURTHER INFORMATION CONTINUED FROM PCT/ISA/210

- 1) claims 1-14, 21, 22, 25-27 all totally; claim 24 partially.

Isolated polynucleotide comprising Seq.ID:1, homologue sequences and derivates. Vectors and transformed host cells. Process for producing human recombinant CTLA-8 protein. Isolated human CTLA-8 protein as in Seq.ID:2 and fragments. Pharmaceutical compositions and uses in therapy. Antibodies.

- 2) claims 15-17 all totally; claim 24 partially.

Therapeutical uses of the protein as in Seq.ID:4 or fragments.

- 3) claims 18-20 all totally; claim 24 partially.

Therapeutical uses of the protein as in Seq.ID:6 or fragments.

- 4) claims 23 all totally; claim 24 partially.

Therapeutical uses of IL-17 or active fragments.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 96/11889

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9518826 A	13-07-95	AU 1520895 A EP 0733069 A JP 9501572 T	01-08-95 25-09-96 18-02-97
WO 9707198 A	27-02-97	AU 6712396 A AU 6768596 A WO 9704097 A	18-02-97 12-03-97 06-02-97
WO 9629408 A	26-09-96	AU 5526396 A	08-10-96



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : C12N 15/19, 15/12, 15/24, C07K 14/52, 14/725, 14/54, A61K 38/17, 38/19, 38/20, C12N 5/10, C07K 16/24, 16/28, A61K 48/00</p>	<p><b>A3</b></p>	<p>(11) International Publication Number: <b>WO 97/04097</b></p> <p>(43) International Publication Date: 6 February 1997 (06.02.97)</p>
<p>(21) International Application Number: PCT/US96/11889</p> <p>(22) International Filing Date: 18 July 1996 (18.07.96)</p> <p>(30) Priority Data: 08/504,032 19 July 1995 (19.07.95) US 08/514,014 11 August 1995 (11.08.95) US</p> <p>(71) Applicant: GENETICS INSTITUTE, INC. [US/US]; 87 Cambridge Park Drive, Cambridge, MA 02140 (US).</p> <p>(72) Inventors: JACOBS, Kenneth; 151 Beaumont Avenue, Newton, MA 02160 (US). KELLEHER, Kerry; 50 Hurley Circle, Marlborough, MA 01752 (US). CARLIN, McKeough; 16 Chauncy Street #2, Cambridge, MA 02138 (US). GOLDMAN, Samuel; 9 Mohawk Drive, Acton, MA 01720 (US). PITTMAN, Debra; 20 N. Shore Road, Windham, NH 03087 (US). MI, Sha; 4 Vernon Road, Belmont, MA 02178 (US). NEBEN, Steven; 13 Duggan Road, Acton, MA 01720 (US). GIANNOTTI, JoAnn; 409 Arlington Street, Acton, MA 01720 (US). GOLDEN-FLEET, Margaret; 19 Usher Road, Medford, MA 02155 (US).</p> <p>(74) Agent: BROWN, Scott, A.; Genetics Institute, Inc., 87 CambridgePark Drive, Cambridge, MA 02140 (US).</p>	<p>(81) Designated States: AU, CA, JP, MX, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p><b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p> <p>(88) Date of publication of the international search report: 12 September 1997 (12.09.97)</p>	
<p>(54) Title: HUMAN CTLA-8 AND USES OF CTLA-8-RELATED PROTEINS</p>		
<p>(57) Abstract</p> <p>Polynucleotides encoding human CTLA-8 and related proteins are disclosed. Human CTLA-8 proteins and methods for their production are also disclosed. Methods of treatment using human CTLA-8 proteins, rat CTLA-8 proteins and herpesvirus herpes CTLA-8 proteins are also provided.</p>		



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

## HUMAN CTLA-8 AND USES OF CTLA-8-RELATED PROTEINS

This application is a continuation-in-part of application Ser. No. 08/504,032, filed July 19, 1995, and a continuation-in-part of application Ser. No. 08/514,014, filed August 11, 1995.

Field of the Invention

5           The present invention relates to human CTLA-8 proteins, nucleic acids encoding such proteins, methods of treatment using such proteins. The invention also relates to the use of rat CTLA-8 proteins and herpesvirus *Saimiri* ORF13 proteins in methods of treatment.

10           Background of the Invention

Cytokines are secreted proteins which act on specific hematopoietic target cells to cause a differentiation event or on other target cells to induce a particular physiological response, such as secretion of proteins characteristic of inflammation. Cytokines, also variously known as lymphokines, hematopoietins, interleukins, colony  
15           stimulating factors, and the like, can be important therapeutic agents, especially for diseases or conditions in which a specific cell population is depleted. For example, erythropoietin, G-CSF, and GM-CSF, have all become important for treatment of anemia and leukopenia, respectively. Other cytokines such as interleukin-3, interleukin-6, interleukin-11 and interleukin-12 show promise in treatment of  
20           conditions such as thrombocytopenia and modulation of immune response.

For these reasons a significant research effort has been expended in searching for novel cytokines and cloning the DNAs which encode them. In the past, novel cytokines were identified by assaying a particular cell such as a bone marrow cell, for a measurable response, such as proliferation. The search for novel cytokines has thus  
5           been limited by the assays available, and if a novel cytokine has an activity which is unmeasurable by a known assay, the cytokine remains undetectable. In a newer approach, cDNAs encoding cytokines have been detected using the polymerase chain

reaction (PCR) and oligonucleotide primers having homology to shared motifs of known cytokines or their receptors. The PCR approach is also limited by the necessity for knowledge of previously cloned cytokines in the same protein family. Cytokines have also been cloned using subtractive hybridization to construct and screen cDNA libraries, or they can potentially be cloned using PCR followed by gel electrophoresis to detect differentially expressed genes. The subtractive hybridization methods are based on the assumption that cytokine mRNAs are those that are differentially expressed, and these methods do not require any prior knowledge of the sequence of interest. However, many cytokines may be encoded by mRNAs which are not differentially expressed, and thus are undetectable using these methods.

It would be desirable to develop new methods for identifying novel cytokines and other secreted factors and to isolate polynucleotides encoding them.

#### Summary of the Invention

In developing the present invention, methods were employed which selectively identify polynucleotides which encode secreted proteins. One such polynucleotide was isolated which encodes "human CTLA-8." In accordance with the present invention, polynucleotides encoding human CTLA-8 and active fragments thereof are disclosed. "CTLA-8" is used throughout the present specification to refer to both proteins and polynucleotides encoding those proteins and to refer to proteins and polynucleotides from all mammalian species.

In certain embodiments, the present invention provides an isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) the nucleotide sequence of SEQ ID NO:1 from nucleotide 146 to nucleotide 544;
- (b) a nucleotide sequence capable of hybridizing to a nucleic acid sequence specified in (a);
- (c) a nucleotide sequence varying from the sequence of the nucleotide sequence specified in (a) as a result of degeneracy of the genetic code; and
- (d) an allelic variant of the nucleotide sequence specified in (a).

Preferably, the polynucleotide of the invention encodes a protein having CTLA-8 activity. In other embodiments the polynucleotide is operably linked to an expression control sequence. In other preferred embodiments, the polynucleotide is contained in a vector suitable for *in vivo* expression in a mammalian subject. Polynucleotides comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 55 to nucleotide 544, the nucleotide sequence of SEQ ID NO:1 from nucleotide 139 to nucleotide 544 or the nucleotide sequence of SEQ ID NO:1 from nucleotide 86 to nucleotide 544 are particularly preferred.

Host cells transformed with the polynucleotides of the invention are also provided, including mammalian cells.

Processes are also provided for producing a human CTLA-8 protein, said processes comprising:

(a) growing a culture of the host cell of the invention in a suitable culture medium; and

(b) purifying the human CTLA-8 protein from the culture.

Isolated human CTLA-8 protein is also provided which comprising an amino acid sequence selected from the group consisting of:

(a) the amino acid sequence of SEQ ID NO:2;

(b) the amino acid sequence of SEQ ID NO:2 from amino acids 11 to 163;

(c) the amino acid sequence of SEQ ID NO:2 from amino acids 29 to 163;

(d) the amino acid sequence of SEQ ID NO:2 from amino acids 31 to 163;

and

(e) fragments of (a), (b), (c) or (d) having CTLA-8 activity.

Proteins comprising the amino acid sequence of SEQ ID NO:2 and comprising the sequence from amino acids 29 to 163, from amino acid 31 to 163, or from amino acids 11 to 163 of SEQ ID NO:2 are particularly preferred. Preferably, the protein has CTLA-8 activity. Pharmaceuticals composition comprising a human CTLA-8 protein of the invention and a pharmaceutically acceptable carrier are also provided.

Compositions are also disclosed which comprise an antibody which specifically reacts with a human CTLA-8 protein of the invention.

Methods of treating a mammalian subject are also provided which comprise administering a therapeutically effective amount of a pharmaceutical composition comprising a human CTLA-8 protein.

5 Rat CTLA-8 and active (*i.e.*, having CTLA-8 activity) fragments thereof may also be used in such methods of treatment. Preferably the rat protein is administered as a composition comprising a pharmaceutically acceptable carrier and a protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:4;
- (b) the amino acid sequence of SEQ ID NO:4 from amino acids 18 to 150;
- 10 and
- (c) fragments of (a) or (b) having CTLA-8 activity.

Herpesvirus *Saimiri* ORF13, referred to herein as "herpes CTLA-8", and active (*i.e.*, having CTLA-8 activity) fragments thereof and active fragments thereof may also be used in such methods of treatment. Preferably the herpes CTLA-8 protein is administered as a composition comprising a pharmaceutically acceptable carrier and

15 a protein comprising an amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence of SEQ ID NO:6;
- (b) the amino acid sequence of SEQ ID NO:6 from amino acids 19 to 151;
- and
- 20 (c) fragments of (a) or (b) having CTLA-8 activity.

The invention also provides a method of treating a mammalian subject comprising administering a therapeutically effective amount of a composition comprising a pharmaceutically acceptable carrier and IL-17 or an active fragment thereof.

25 In methods of treatment provided by the present invention, preferably the subject is treated to produce an effect selected from the group consisting of inhibition of angiogenesis, inhibition of growth or proliferation of vascular endothelial cells, inhibition of tumor growth, inhibition of angiogenesis-dependent tissue growth, proliferation of myeloid cells or progenitors, proliferation of erythroid cells or

30 progenitors, proliferation of lymphoid cells or progenitors, induction of IFN $\gamma$  production, induction of IL-3 production and induction of GM-CSF production.

### Brief Description of the Figures

Fig. 1 is a comparison of homologous regions of the amino acid sequences of human CTLA-8 (indicated as "B18\_F1"), rat CTLA-8 (indicated as "Musctla8") and herpes CTLA-8 (indicated as "Hsvie\_2").

5        Fig. 2 depicts autoradiographs demonstrating expression of human CTLA-8 in COS cells.

Fig. 3 presents data relating to the ability of human CTLA-8 to inhibit angiogenesis.

10       Figs. 4 and 5 present data relating to the ability of human CTLA-8 to produce or induce hematopoietic activity.

Figs. 6 and 7 present data demonstrating the ability of human CTLA-8 to induce production of IL-6 and IL-8.

### Detailed Description of Preferred Embodiments

15       The inventors of the present application have identified and provided a polynucleotide encoding a human CTLA-8 protein. SEQ ID NO:1 provides the nucleotide sequence of a cDNA encoding the human CTLA-8 protein. SEQ ID NO:2 provides the amino acid sequence of the human CTLA-8 protein. Alternatively, the initiating methionine may be at amino acid 11 of SEQ ID NO:2. On the basis of  
20       amino terminal sequencing, the mature protein sequence is believed to begin at amino acid 31 of SEQ ID NO:2 (encoded by the sequence beginning with nucleotide 146 of SEQ ID NO:1).

25       The region from amino acid 29 to amino acid 163 of human CTLA-8 (SEQ ID NO:2) shows marked homology to portions of rat CTLA-8 (amino acids 18 to 150 of SEQ ID NO:4) and herpesvirus *Saimiri* ORF13 ("herpes CTLA-8") (amino acids 19 to 151 of SEQ ID NO:5). A cDNA sequence encoding rat CTLA-8 is listed at SEQ ID NO:3 and its corresponding amino acid sequence is reported at SEQ ID NO:4. A cDNA sequence encoding herpes CTLA-8 is listed at SEQ ID NO:5 and its corresponding amino acid sequence is reported at SEQ ID NO:6. Homology between  
30       rat CTLA-8 and herpes CTLA-8 was reported by Rouvier et al., J. Immunol. 1993, 150, 5445-5456.

Applicants had previously incorrectly identified the rat sequences of SEQ ID NO:3 and SEQ ID NO:4 as applying to murine CTLA-8. Applicants' human CTLA-8 (B18) does also show homolgy to the true murine CTLA-8 sequence.

5       Golstein et al. (WO95/18826; Fossiez et al., Microbial Evasion and Subversion of Immunity 544:3222 (Abstract)) have also reported a species they initially identified as "human CTLA-8." However, examination of the sequence of the Golstein et al. species and the human CTLA-8 (B18) sequence of the present invention readily reveals that they are two different proteins, although they are homologous with each other and with the rat CTLA-8 and herpes CTLA-8 identified herein. The Golstein  
10       et al. species has now been renamed as interleukin-17 (IL-17). Because of the homology between applicants' human CTLA-8 (B18) and IL-17, these proteins are expected to share some activities.

It has also been preliminarily determined that human CTLA-8 (B18) forms homodimers when expressed. As a result, human CTLA-8 proteins may possess  
15       activity in either monomeric or dimeric forms. Human CTLA-8 proteins can also be produced as heterodimers with rat and herpes CTLA-8 proteins and with human IL-17. These heterodimers are also expected to have activities of the proteins of which they are comprised.

Forms of human CTLA-8 protein of less than full length are encompassed  
20       within the present invention and may be produced by expressing a corresponding fragment of the polynucleotide encoding the human CTLA-8 protein (SEQ ID NO:1). These corresponding polynucleotide fragments are also part of the present invention. Modified polynucleotides as described above may be made by standard molecular biology techniques, including site-directed mutagenesis methods which are known in  
25       the art or by the polymerase chain reaction using appropriate oligonucleotide primers.

For the purposes of the present invention, a protein has "CTLA-8 activity" if it either (1) displays biological activity in a factor-dependent cell proliferation assay (preferably an assay in which full-length the corresponding species full-length CTLA-8 is active) (including without limitation those assays described below), or (2) induces  
30       expression or secretion of  $\gamma$ -IFN, or (3) displays chemoattractant or chemotactic activity in a chemoattraction or chemotaxis assay (preferably as assay in which full-

length the corresponding species full-length CTLA-8 is active) or (4) induces expression of secretion of IL-3 or GM-CSF.

Human CTLA-8 protein or fragments thereof having CTLA-8 activity may be fused to carrier molecules such as immunoglobulins. For example, human CTLA-8 protein may be fused through "linker" sequences to the Fc portion of an immunoglobulin.

The invention also encompasses allelic variations of the nucleotide sequence as set forth in SEQ ID NO:1, that is, naturally-occurring alternative forms of the isolated polynucleotide of SEQ ID NO:1 which also encode human CTLA-8 or CTLA-8 proteins having CTLA-8 activity. Also included in the invention are isolated polynucleotides which hybridize to the nucleotide sequence set forth in SEQ ID NO:1 under highly stringent (0.2xSSC at 65°C), stringent (e.g. 4xSSC at 65 C or 50% formamide and 4xSSC at 42°C), or relaxed (4xSSC at 50°C or 30-40% formamide and 4xSSC at 42°C) conditions. Isolated polynucleotides which encode human CTLA-8 protein but which differ from the nucleotide sequence set forth in SEQ ID NO:1 by virtue of the degeneracy of the genetic code are also encompassed by the present invention. Variations in the nucleotide sequence as set forth in SEQ ID NO:1 which are caused by point mutations or by induced modifications which enhance CTLA-8 activity, half-life or production level are also included in the invention.

The isolated polynucleotides of the invention may be operably linked to an expression control sequence such as the pMT2 or pED expression vectors disclosed in Kaufman et al., Nucleic Acids Res. 19, 4485-4490 (1991), in order to produce the CTLA-8 protein recombinantly. Many suitable expression control sequences are known in the art. General methods of expressing recombinant proteins are also known and are exemplified in R. Kaufman, Methods in Enzymology 185, 537-566 (1990). As defined herein "operably linked" means enzymatically or chemically ligated to form a covalent bond between the isolated polynucleotide of the invention and the expression control sequence, in such a way that the CTLA-8 protein is expressed by a host cell which has been transformed (transfected) with the ligated polynucleotide/expression control sequence.



A number of types of cells may act as suitable host cells for expression of the human CTLA-8 protein. Any cell type capable of expressing functional human CTLA-8 protein may be used. Suitable mammalian host cells include, for example, monkey COS cells, Chinese Hamster Ovary (CHO) cells, human kidney 293 cells, human epidermal A431 cells, human Colo205 cells, 3T3 cells, CV-1 cells, other  
5 transformed primate cell lines, normal diploid cells, cell strains derived from *in vitro* culture of primary tissue, primary explants, HeLa cells, mouse L cells, BHK, HL-60, U937, HaK, Rat2, BaF3, 32D, FDCP-1, PC12 or C2C12 cells.

The human CTLA-8 protein may also be produced by operably linking the  
10 isolated polynucleotide of the invention to suitable control sequences in one or more insect expression vectors, and employing an insect expression system. Materials and methods for baculovirus/insect cell expression systems are commercially available in kit form from, e.g., Invitrogen, San Diego, California, U.S.A. (the MaxBac® kit), and such methods are well known in the art, as described in Summers and Smith, Texas  
15 Agricultural Experiment Station Bulletin No. 1555 (1987), incorporated herein by reference. Soluble forms of the human CTLA-8 protein may also be produced in insect cells using appropriate isolated polynucleotides as described above.

Alternatively, the human CTLA-8 protein may be produced in lower eukaryotes such as yeast or in prokaryotes such as bacteria. Suitable yeast strains  
20 include *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces* strains, *Candida*, or any yeast strain capable of expressing heterologous proteins. Suitable bacterial strains include *Escherichia coli*, *Bacillus subtilis*, *Salmonella typhimurium*, or any bacterial strain capable of expressing heterologous proteins.

The human CTLA-8 protein of the invention may also be expressed as a  
25 product of transgenic animals, e.g., as a component of the milk of transgenic cows, goats, pigs, or sheep which are characterized by somatic or germ cells containing a polynucleotide sequence encoding the human CTLA-8 protein.

The human CTLA-8 protein of the invention may be prepared by growing a culture of transformed host cells under culture conditions necessary to express the  
30 desired protein. The resulting expressed protein may then be purified from the culture medium or cell extracts. Soluble forms of the human CTLA-8 protein of the invention

can be purified from conditioned media. Membrane-bound forms of human CTLA-8 protein of the invention can be purified by preparing a total membrane fraction from the expressing cell and extracting the membranes with a non-ionic detergent such as Triton X-100.

5           The human CTLA-8 protein can be purified using methods known to those skilled in the art. For example, the human CTLA-8 protein of the invention can be concentrated using a commercially available protein concentration filter, for example, an Amicon or Millipore Pellicon ultrafiltration unit. Following the concentration step, the concentrate can be applied to a purification matrix such as a gel filtration medium.  
10           Alternatively, an anion exchange resin can be employed; for example, a matrix or substrate having pendant diethylaminoethyl (DEAE) groups. The matrices can be acrylamide, agarose, dextran, cellulose or other types commonly employed in protein purification. Alternatively, a cation exchange step can be employed. Suitable cation exchangers include various insoluble matrices comprising sulfopropyl or  
15           carboxymethyl groups. Sulfopropyl groups are preferred (e.g., S-Sepharose® columns). The purification of the human CTLA-8 protein from culture supernatant may also include one or more column steps over such affinity resins as concanavalin A-agarose, heparin-toyopearl® or Cibacrom blue 3GA Sepharose®; or by hydrophobic interaction chromatography using such resins as phenyl ether, butyl ether,  
20           or propyl ether; or by immunoaffinity chromatography. Finally, one or more reverse-phase high performance liquid chromatography (RP-HPLC) steps employing hydrophobic RP-HPLC media, e.g., silica gel having pendant methyl or other aliphatic groups, can be employed to further purify the human CTLA-8 protein. Some or all of the foregoing purification steps, in various combinations or with other known  
25           methods, can also be employed to provide a substantially purified isolated recombinant protein.

          Preferably, the human CTLA-8 protein is purified so that it is substantially free of other mammalian proteins.

          It is believed that human CTLA-8, active fragments and variants thereof, and  
30           CTLA-8 related proteins (such as, for example, rat CTLA-8 and herpes CTLA-8) (collectively "CTLA-8 proteins") possess or induce cytokine activities. Human

CTLA-8 expression correlated with  $\gamma$ -IFN expression in induced primary cells and can induce the expression of IL-3 and/or GM-CSF, which expression can in turn produce effects associated with the induced cytokine. Therefore, human CTLA-8 and CTLA-8 related proteins may have an effect on proliferation or function of myeloid cells, erythroid cells, lymphoid cells and their progenitors. Human CTLA-8 proteins may also play a role in formation of platelets or their progenitors.

A protein of the present invention may exhibit cytokine, cell proliferation (either inducing or inhibiting) or cell differentiation (either inducing or inhibiting) activity or may induce production of other cytokines in certain cell populations. Many protein factors discovered to date, including all known cytokines, have exhibited activity in one or more factor dependent cell proliferation assays, and hence the assays serve as a convenient confirmation of cytokine activity. The activity of a protein of the present invention is evidenced by any one of a number of routine factor dependent cell proliferation assays for cell lines including, without limitation, 32D, DA2, DA1G, T10, B9, B9/11, BaF3, MC9/G, M+ (preB M+), 2E8, RB5, DA1, 123, T1165, HT2, CTLL2, TF-1, Mo7e and CMK.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Assays for T-cell or thymocyte proliferation include without limitation those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., J. Immunol. 137:3494-3500, 1986; Bertagnolli et al., J. Immunol. 145:1706-1712, 1990; Bertagnolli et al., Cellular Immunology 133:327-341, 1991; Bertagnolli, et al., J. Immunol. 149:3778-3783, 1992; Bowman et al., J. Immunol. 152: 1756-1761, 1994.

Assays for cytokine production and/or proliferation of spleen cells, lymph node cells or thymocytes include, without limitation, those described in: Polyclonal T cell stimulation, Kruisbeek, A.M. and Shevach, E.M. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 3.12.1-3.12.14, John Wiley and Sons, Toronto. 1994; and Measurement of mouse and human Interferon  $\gamma$ , Schreiber, R.D.

In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.8.1-6.8.8, John Wiley and Sons, Toronto. 1994.

Assays for proliferation and differentiation of hematopoietic and lymphopoietic cells include, without limitation, those described in: Measurement of Human and Murine Interleukin 2 and Interleukin 4, Bottomly, K., Davis, L.S. and Lipsky, P.E. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.3.1-6.3.12, John Wiley and Sons, Toronto. 1991; deVries et al., J. Exp. Med. 173:1205-1211, 1991; Moreau et al., Nature 336:690-692, 1988; Greenberger et al., Proc. Natl. Acad. Sci. U.S.A. 80:2931-2938, 1983; Measurement of mouse and human interleukin 6 - Nordan, R. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.6.1-6.6.5, John Wiley and Sons, Toronto. 1991; Smith et al., Proc. Natl. Acad. Sci. U.S.A. 83:1857-1861, 1986; Measurement of human Interleukin 11 - Bennett, F., Giannotti, J., Clark, S.C. and Turner, K. J. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.15.1 John Wiley and Sons, Toronto. 1991; Measurement of mouse and human Interleukin 9 - Ciarletta, A., Giannotti, J., Clark, S.C. and Turner, K.J. In *Current Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 6.13.1, John Wiley and Sons, Toronto. 1991.

Assays for T-cell clone responses to antigens (which will identify, among others, proteins that affect APC-T cell interactions as well as direct T-cell effects by measuring proliferation and cytokine production) include, without limitation, those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober  
Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function; Chapter 6, Cytokines and their cellular receptors; Chapter 7, Immunologic studies in Humans); Weinberger et al., Proc. Natl. Acad. Sci. USA 77:6091-6095, 1980; Weinberger et al., Eur. J. Immun. 11:405-411, 1981; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988.

A protein of the present invention may also exhibit immune stimulating or immune suppressing activity, including without limitation the activities for which assays are described herein. A protein may be useful in the treatment of various

immune deficiencies and disorders (including severe combined immunodeficiency (SCID)), e.g., in regulating (up or down) growth and proliferation of T and/or B lymphocytes, as well as effecting the cytolytic activity of NK cells and other cell populations. These immune deficiencies may be genetic or be caused by viral (e.g., HIV) as well as bacterial or fungal infections, or may result from autoimmune disorders. More specifically, infectious diseases caused by viral, bacterial, fungal or other infection may be treatable using a protein of the present invention, including infections by HIV, hepatitis viruses, herpes viruses, mycobacteria, leishmania, malaria and various fungal infections such as candida. Of course, in this regard, a protein of the present invention may also be useful where a boost to the immune system generally would be indicated, i.e., in the treatment of cancer.

Autoimmune disorders which may be treated using a protein of the present invention include, for example, multiple sclerosis, systemic lupus erythematosus, rheumatoid arthritis, autoimmune pulmonary inflammation, Guillain-Barre syndrome, autoimmune thyroiditis, insulin dependent diabetes mellitus, myasthenia gravis, graft-versus-host disease and autoimmune inflammatory eye disease. Such a protein of the present invention may also be useful in the treatment of allergic reactions and conditions, such as asthma or other respiratory problems. Other conditions, in which immune suppression is desired (including, for example, asthma and related respiratory conditions), may also be treatable using a protein of the present invention.

A protein of the present invention may also suppress chronic or acute inflammation, such as, for example, that associated with infection (such as septic shock or systemic inflammatory response syndrome (SIRS)), inflammatory bowel disease, Crohn's disease or resulting from over production of cytokines such as TNF or IL-1 (such as the effect demonstrated by IL-11).

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for thymocyte or splenocyte cytotoxicity include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte

- Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al., J. Immunol. 135:1564-1572, 1985; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988;
- 5 Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al., J. Immunol. 135:1564-1572, 1985; Takai et al., J. Immunol. 137:3494-3500, 1986; Bowman et al., J. Virology 61:1992-1998; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnolli et al., Cellular Immunology 133:327-341, 1991; Brown et al., J. Immunol. 153:3079-3092, 1994.
- 10 Assays for T-cell-dependent immunoglobulin responses and isotype switching (which will identify, among others, proteins that modulate T-cell dependent antibody responses and that affect Th1/Th2 profiles) include, without limitation, those described in: Maliszewski, J. Immunol. 144:3028-3033, 1990; and Assays for B cell function: *In vitro* antibody production, Mond, J.J. and Brunswick, M. In *Current*
- 15 *Protocols in Immunology*. J.E.e.a. Coligan eds. Vol 1 pp. 3.8.1-3.8.16, John Wiley and Sons, Toronto. 1994.
- Mixed lymphocyte reaction (MLR) assays (which will identify, among others, proteins that generate predominantly Th1 and CTL responses) include, without limitation, those described in: *Current Protocols in Immunology*, Ed by J. E. Coligan,
- 20 A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, *In Vitro* assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnolli et al., J. Immunol. 149:3778-3783, 1992.
- 25 Dendritic cell-dependent assays (which will identify, among others, proteins expressed by dendritic cells that activate naive T-cells) include, without limitation, those described in: Guery et al., J. Immunol. 134:536-544, 1995; Inaba et al., *Journal of Experimental Medicine* 173:549-559, 1991; Macatonia et al., *Journal of Immunology* 154:5071-5079, 1995; Porgador et al., *Journal of Experimental Medicine*
- 30 182:255-260, 1995; Nair et al., *Journal of Virology* 67:4062-4069, 1993; Huang et al., *Science* 264:961-965, 1994; Macatonia et al., *Journal of Experimental Medicine*

169:1255-1264, 1989; Bhardwaj et al., Journal of Clinical Investigation 94:797-807, 1994; and Inaba et al., Journal of Experimental Medicine 172:631-640, 1990.

Assays for lymphocyte survival/apoptosis (which will identify, among others, proteins that prevent apoptosis after superantigen induction and proteins that regulate lymphocyte homeostasis) include, without limitation, those described in:

5 Darzynkiewicz et al., Cytometry 13:795-808, 1992; Gorczyca et al., Leukemia 7:659-670, 1993; Gorczyca et al., Cancer Research 53:1945-1951, 1993; Itoh et al., Cell 66:233-243, 1991; Zacharchuk, Journal of Immunology 145:4037-4045, 1990; Zamai et al., Cytometry 14:891-897, 1993; Gorczyca et al., International Journal of

10 Oncology 1:639-648, 1992.

Assays for proteins that influence early steps of T-cell commitment and development include, without limitation, those described in: Antica et al., Blood 84:111-117, 1994; Fine et al., Cellular Immunology 155:111-122, 1994; Galy et al., Blood 85:2770-2778, 1995; Toki et al., Proc. Nat. Acad Sci. USA 88:7548-7551,

15 1991.

A protein of the present invention may be useful in regulation of hematopoiesis and, consequently, in the treatment of myeloid or lymphoid cell deficiencies. Even marginal biological activity in support of colony forming cells or of factor-dependent cell lines indicates involvement in regulating hematopoiesis, e.g. in supporting the

20 growth and proliferation of erythroid progenitor cells alone or in combination with other cytokines, thereby indicating utility, for example, in treating various anemias or for use in conjunction with irradiation/chemotherapy to stimulate the production of erythroid precursors and/or erythroid cells; in supporting the growth and proliferation of myeloid cells such as granulocytes and monocytes/macrophages (i.e., traditional

25 CSF activity) useful, for example, in conjunction with chemotherapy to prevent or treat consequent myelo-suppression; in supporting the growth and proliferation of megakaryocytes and consequently of platelets thereby allowing prevention or treatment of various platelet disorders such as thrombocytopenia, and generally for use in place of or complementarily to platelet transfusions; and/or in supporting the growth

30 and proliferation of hematopoietic stem cells which are capable of maturing to any and all of the above-mentioned hematopoietic cells and therefore find therapeutic utility.

in various stem cell disorders (such as those usually treated with transplantation, including, without limitation, aplastic anemia and paroxysmal nocturnal hemoglobinuria), as well as in repopulating the stem cell compartment post irradiation/chemotherapy, either *in-vivo* or *ex-vivo* (i.e. in conjunction with bone marrow transplantation) as normal cells or genetically manipulated for gene therapy.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for proliferation and differentiation of various hematopoietic lines are cited above.

Assays for embryonic stem cell differentiation (which will identify, among others, proteins that influence embryonic differentiation hematopoiesis) include, without limitation, those described in: Johansson et al. *Cellular Biology* 15:141-151, 1995; Keller et al., *Molecular and Cellular Biology* 13:473-486, 1993; McClanahan et al., *Blood* 81:2903-2915, 1993.

Assays for stem cell survival and differentiation (which will identify, among others, proteins that regulate lympho-hematopoiesis) include, without limitation, those described in: Methylcellulose colony forming assays, Freshney, M.G. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 265-268, Wiley-Liss, Inc., New York, NY. 1994; Hirayama et al., *Proc. Natl. Acad. Sci. USA* 89:5907-5911, 1992; Primitive hematopoietic colony forming cells with high proliferative potential, McNiece, I.K. and Briddell, R.A. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 23-39, Wiley-Liss, Inc., New York, NY. 1994; Neben et al., *Experimental Hematology* 22:353-359, 1994; Cobblestone area forming cell assay, Ploemacher, R.E. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 1-21, Wiley-Liss, Inc., New York, NY. 1994; Long term bone marrow cultures in the presence of stromal cells, Spooncer, E., Dexter, M. and Allen, T. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 163-179, Wiley-Liss, Inc., New York, NY. 1994; Long term culture initiating cell assay, Sutherland, H.J. In *Culture of Hematopoietic Cells*. R.I. Freshney, et al. eds. Vol pp. 139-162, Wiley-Liss, Inc., New York, NY. 1994.



CTLA-8 proteins are useful in the treatment of various immune deficiencies and disorders (including SCID), *e.g.*, in regulating (up or down) growth, proliferation and/or activity of T and/or B lymphocytes, as well as the cytolytic activity of NK cells. These immune deficiencies may be caused by viral (*e.g.*, HIV) as well as bacterial infections, or may result from autoimmune disorders. More specifically, infectious diseases caused by viral, bacterial, fungal or other infection may be treatable using CTLA-8 proteins, including infections by HIV, hepatitis, influenza, CMV, herpes, mycobacterium, leishmaniasis, malaria and various fungal infections (such as candida). Of course, in this regard, the CTLA-8 proteins may also be useful where a boost to the immune system generally would be indicated, *i.e.*, in the treatment of cancer or as an adjuvant to vaccines. Autoimmune disorders which may be treated using factors of the present invention include, for example, multiple sclerosis, systemic lupus erythematosus, rheumatoid arthritis, autoimmune pulmonary inflammation, Guillain-Barre syndrome, autoimmune thyroiditis, insulin dependent diabetes melitis and autoimmune inflammatory eye disease. The CTLA-8 proteins are also expected to be useful in the treatment of allergic reactions and conditions.

CTLA-8 proteins are also expected to have chemotactic activity. A protein or peptide has "chemotactic activity," as used herein, if it can stimulate, directly or indirectly, the directed orientation or movement of cells, including myeloid and lymphoid cells. Preferably, the protein or peptide has the ability to directly stimulate directed movement of cells (particularly T-cells). Whether a particular protein or peptide has chemotactic activity for cells can be readily determined by employing such protein or peptide in any known assay for cell chemotaxis.

CTLA-8 proteins also inhibit growth and proliferation of vascular endothelial cells. As a result, human CTLA-8 proteins are effective in inhibiting angiogenesis (*i.e.*, vascular formation). This activity will also be useful in the treatment of tumors and other conditions in which angiogenesis is involved. Inhibition of angiogenesis by human CTLA-8 proteins will also result in inhibition or prevention of the condition to which normal angiogenesis would contribute.

Isolated CTLA-8 proteins, purified from cells or recombinantly produced, may be used as a pharmaceutical composition when combined with a pharmaceutically

acceptable carrier. Such a composition may contain, in addition to CTLA-8 protein and carrier, diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredient(s). The characteristics of the carrier will depend on the route of administration. The pharmaceutical composition of the invention may also contain cytokines, lymphokines, or other hematopoietic factors such as M-CSF, GM-CSF, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-15, G-CSF,  $\gamma$ -IFN, stem cell factor, and erythropoietin. The pharmaceutical composition may contain thrombolytic or anti-thrombotic factors such as plasminogen activator and Factor VIII. The pharmaceutical composition may further contain other anti-inflammatory agents. Such additional factors and/or agents may be included in the pharmaceutical composition to produce a synergistic effect with CTLA-8 protein, or to minimize side effects caused by the CTLA-8 protein. Conversely, CTLA-8 protein may be included in formulations of the particular cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent to minimize side effects of the cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent.

The pharmaceutical composition of the invention may be in the form of a liposome in which CTLA-8 protein is combined, in addition to other pharmaceutically acceptable carriers, with amphipathic agents such as lipids which exist in aggregated form as micelles, insoluble monolayers, liquid crystals, or lamellar layers which in aqueous solution. Suitable lipids for liposomal formulation include, without limitation, monoglycerides, diglycerides, sulfatides, lysolecithin, phospholipids, saponin, bile acids, and the like. Preparation of such liposomal formulations is within the level of skill in the art, as disclosed, for example, in U.S. Patent No. 4,235,871; U.S. Patent No. 4,501,728; U.S. Patent No. 4,837,028; and U.S. Patent No. 4,737,323, all of which are incorporated herein by reference.

As used herein, the term "therapeutically effective amount" means the total amount of each active component of the pharmaceutical composition or method that is sufficient to show a meaningful patient benefit, e.g., amelioration of symptoms of,

healing of, or increase in rate of healing of such conditions. When applied to an individual active ingredient, administered alone, the term refers to that ingredient alone. When applied to a combination, the term refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in  
5 combination, serially or simultaneously.

In practicing the method of treatment or use of the present invention, a therapeutically effective amount of CTLA-8 protein is administered to a mammal. CTLA-8 protein may be administered in accordance with the method of the invention either alone or in combination with other therapies such as treatments employing  
10 cytokines, lymphokines or other hematopoietic factors. When co-administered with one or more cytokines, lymphokines, other hematopoietic factors or vaccine components (such as antigens or other adjuvants), CTLA-8 protein may be administered either simultaneously with the cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors, or sequentially. If  
15 administered sequentially, the attending physician will decide on the appropriate sequence of administering CTLA-8 protein in combination with cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors.

Administration of CTLA-8 protein used in the pharmaceutical composition or to practice the method of the present invention can be carried out in a variety of  
20 conventional ways, such as oral ingestion, inhalation, or cutaneous, subcutaneous, or intravenous injection. Intravenous administration to the patient is preferred.

When a therapeutically effective amount of CTLA-8 protein is administered orally, CTLA-8 protein will be in the form of a tablet, capsule, powder, solution or elixir. When administered in tablet form, the pharmaceutical composition of the  
25 invention may additionally contain a solid carrier such as a gelatin or an adjuvant. The tablet, capsule, and powder contain from about 5 to 95% CTLA-8 protein, and preferably from about 25 to 90% CTLA-8 protein. When administered in liquid form, a liquid carrier such as water, petroleum, oils of animal or plant origin such as peanut oil, mineral oil, soybean oil, or sesame oil, or synthetic oils may be added. The liquid  
30 form of the pharmaceutical composition may further contain physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol.

propylene glycol or polyethylene glycol. When administered in liquid form, the pharmaceutical composition contains from about 0.5 to 90% by weight of CTLA-8 protein, and preferably from about 1 to 50% CTLA-8 protein.

5 When a therapeutically effective amount of CTLA-8 protein is administered by intravenous, cutaneous or subcutaneous injection, CTLA-8 protein will be in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable protein solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred pharmaceutical composition for intravenous, cutaneous, or subcutaneous injection should contain, in  
10 addition to CTLA-8 protein an isotonic vehicle such as Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, Lactated Ringer's Injection, or other vehicle as known in the art. The pharmaceutical composition of the present invention may also contain stabilizers, preservatives, buffers, antioxidants, or other additive known to those of skill in the art.

15 The amount of CTLA-8 protein in the pharmaceutical composition of the present invention will depend upon the nature and severity of the condition being treated, and on the nature of prior treatments which the patient has undergone. Ultimately, the attending physician will decide the amount of CTLA-8 protein with which to treat each individual patient. Initially, the attending physician will  
20 administer low doses of CTLA-8 protein and observe the patient's response. Larger doses of CTLA-8 protein may be administered until the optimal therapeutic effect is obtained for the patient, and at that point the dosage is not generally increased further. It is contemplated that the various pharmaceutical compositions used to practice the method of the present invention should contain about 0.1  $\mu$ g to about 100 mg of  
25 CTLA-8 protein per kg body weight, preferably about 0.1  $\mu$ g to about 10 mg of CTLA-8 protein per kg body weight, more preferably about 0.1  $\mu$ g to about 100  $\mu$ g of CTLA-8 protein per kg body weight, most preferably preferably about 0.1  $\mu$ g to about 10  $\mu$ g of CTLA-8 protein per kg body weight.

30 The duration of intravenous therapy using the pharmaceutical composition of the present invention will vary, depending on the severity of the disease being treated and the condition and potential idiosyncratic response of each individual patient. It

is contemplated that the duration of each application of the CTLA-8 protein will be in the range of 12 to 24 hours of continuous intravenous administration. Ultimately the attending physician will decide on the appropriate duration of intravenous therapy using the pharmaceutical composition of the present invention.

5 CTLA-8 protein of the invention may also be used to immunize animals to obtain polyclonal and monoclonal antibodies which specifically react with the CTLA-8 protein and which may inhibit CTLA-8 binding to its receptor. Such antibodies are also useful for performing diagnostics assays for CTLA-8 in accordance with known methods. Such antibodies may be obtained using the entire CTLA-8 protein as an  
10 immunogen, or by using fragments of human CTLA-8 protein. The peptide immunogens additionally may contain a cysteine residue at the carboxyl terminus, and are conjugated to a hapten such as keyhole limpet hemocyanin (KLH). Additional peptide immunogens may be generated by replacing tyrosine residues with sulfated tyrosine residues. Methods for synthesizing such peptides are known in the art, for  
15 example, as in R.P. Merrifield, J.Amer.Chem.Soc. 85, 2149-2154 (1963); J.L. Krstenansky, et al., FEBS Lett. 211, 10 (1987).

Neutralizing or non-neutralizing antibodies (preferably monoclonal antibodies) binding to human CTLA-8 protein may also be useful therapeutics for certain tumors and also in the treatment of conditions described above. These neutralizing  
20 monoclonal antibodies are capable of blocking the ligand binding to the human CTLA-8 protein or mayh promote clearance of protein from the patient.

Because of their homology to human CTLA-8, rat CTLA-8 proteins, herpes CTLA-8 proteins and IL-17 proteins (the "human CTLA-8" of Golstein et al., supra) will also possess CTLA-8 activity as described above. As a result, rat and herpes  
25 CTLA-8 proteins and IL-17 proteins, as well as active fragments and variants thereof, can be used in preparation of pharmaceutical compositions and in methods of treatment as described for human CTLA-8. Rat and herpes CTLA-8 proteins, and active fragments and variants thereof, can be produced as described above using the polynucleotides (or fragments or variants thereof) described in SEQ ID NO:3 and SEQ  
30 ID NO:5, respectively. Rat and herpes CTLA-8 may also be produced as described in Rouvier et al., J. Immunol. 1993, 150, 5445-5456. CTLA-8 proteins of other

species can also be used as described herein. cDNAs encoding rat CTLA-8 and herpes CTLA-8 were deposited with the American Type Culture Collection on July 6, 1995 and assigned accession numbers ATCC 69867 and ATCC 69866, respectively. IL-17 proteins may also be produced as described in Golstein et al., supra.

5           Because of its homology to IL-17, the human CTLA-8 (B18) proteins of the present invention may also share some activities with IL-17.

For the purposes of treatment or therapy, any of the proteins discussed or disclosed herein may be administered by *in vivo* expression of the protein in a mammalian subject. In such instances, a polynucleotide encoding the desired protein is administered to the subject in manner allowing expression in accordance with  
10           known methods, including without limitation the adenovirus methods disclosed herein.

#### Example 1

##### 15           Isolation of Human CTLA-8 cDNA

A partial clone for human CTLA-8 was isolated from a cDNA library made from RNA isolated from stimulated human peripheral blood mononuclear cells. This partial was identified as "B18." B18 is sometimes used herein to refer to the human CTLA-8 of the present invention. Homology searches identified this partial clone as  
20           being related to the herpes and rat CTLA-8 genes. DNA sequence of this partial clone was used to isolate the full-length clone.

In order to isolate a full-length cDNA for B18, a directional, full-length cDNA library by standard means in the COS expression vector pMV2. The cDNA library was transformed into *E. coli* by electroporation. The bulk of the original transformed  
25           cDNA library was frozen in glycerol at -80°C. An aliquot was titered to measure the concentration of transformed *E. coli*. The *E. coli* were thawed, diluted to 76,000/0.1 ml in media containing ampicillin, and 0.1 ml was distributed into the wells of a microtiter dish in an 8 x 8 array. The microtiter dish was placed at 37°C overnight to grow the *E. coli*.

30           To prepare DNA for PCR, 20 µl aliquots of culture from each well were withdrawn and pooled separately for each row and column of eight wells, giving 16

pools of 160  $\mu$ l each. The *E. coli* were pelleted, resuspended in 160  $\mu$ l of standard lysis buffer consisting of 10 mM TrisHCl pH8, 1 mM EDTA, 0.01% Triton X-100, and lysed by heating to 95°C for 10 minutes.

5 To identify which of the wells contained *E. coli* transformed with B18, PCR was performed first on the DNA preps corresponding to the eight columns. The PCR consisted of two sequential reactions with nested oligonucleotides using standard conditions. The oligonucleotides used for the PCR reaction were derived from the sequence of the partial B18 clone. They were:

10 B185: CACAGGCATACACAGGAAGATACATTCA (SEQ ID NO:7)  
B183: TCTTGCTGGATGGGAACGGAATTCA (SEQ ID NO:8)  
B18N: ATACATTCACAGAAGAGCTTCCTGCACA (SEQ ID NO:9)

15 The PCR conditions were 2.5 mM MgCl<sub>2</sub> and 95°C x 2 min for one cycle, 95°C x 1 min plus 68°C x 1 min for 30 cycles, and 68°C x 10 for one cycle. Each reaction was 20  $\mu$ l. The first reaction contained oligonucleotides B185 and B183 and 1  $\mu$ l of the DNA preparations. The second reaction contained oligonucleotides B183 and B18N and 1  $\mu$ l of the first reaction.

20 DNA preps that potentially contained a full-length B18 cDNA clone were identified by agarose gel electrophoresis on an aliquot of the second PCR reaction. A DNA band of the correct mobility was assumed to be derived from a B18 cDNA. Next the same sequence of PCR reactions and gel analysis was done on the DNA preps corresponding to the eight rows. The intersection of a row and a column identified well A2 as potentially containing B18, narrowing it down to the 76,000 *E.*  
25 *coli* originally seeded into that well.

To further purify the individual *E. coli* containing the putative full-length B18 cDNA clone, the concentration of *E. coli* in well A2 was measured by titring and plating dilutions of the well. Then 7600 *E. coli* were seeded into the wells of a second microtiter plate in an 8 x 8 array. The *E. coli* were grown overnight; wells were  
30 pooled, and DNA was prepared as described above. To identify which of these wells contained *E. coli* transformed with B18, sequential PCR reactions were performed

essentially as described above. Agarose gel electrophoresis identified well B2 as potentially containing a B18 cDNA.

5 The *E. coli* containing this cDNA was further purified by seeding wells of a microtiter plate with 253 *E. coli* per well and proceeding as for the purification of the *E. coli* in well A2. Well C3 was identified as containing a putative full-length B18 cDNA clone. The exact *E. coli* was identified by plating the contents of the well onto bacterial culture media and then screening the *E. coli* colonies following established protocols. The probe for these hybridizations was a PCR fragment generated by doing a PCR reaction on the B18 clone using as primers the oligonucleotides described  
10 above (SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9). When a single colony was identified, DNA was prepared and sequenced by standard methods. Comparison of this sequence to the sequence of the original partial clone confirmed identity and that the isolated cDNA was full-length.

15 The full-length clone was deposited with the American Type Culture Collection on July 6, 1995 and assigned accession number ATCC 69868.

### Example 2

#### Expression of Human CTLA-8

20 The full-length B18 clone for human CTLA-8 was transfected into COS cells which were then labelled with <sup>35</sup>S-methionine. An aliquot of conditioned medium from the transfected cell culture was reduced, denatured and electrophoresed on polyacrylamide gels. Autoradiographs of those gels are reproduced in Fig. 2. The band indicated by the arrow demonstrates expression of human CTLA-8.

### Example 3

#### Inhibition of Angiogenesis by Human CTLA-8

25 The ability of human CTLA-8 to inhibit angiogenesis was examined in an angiostatic activity assay (endothelial cell proliferation assay). The assay was done in a 96 well plate. Primary human umbilical cells (HUVECs) were seeded to  $2 \times 10^3$   
30 cells per well in EGM medium (Clonetics)/20% FCS and incubated at 37°C for 24 hr. The cells were then starved in M199 medium (GIBCO BRL) containing 10% charcoal



5 treated serum (M199-CS) for 48 hr at 37°C. Conditioned media containing B18 (human CTLA-8) was obtained from transfected COS or stably expressing CHO cells and 1:10, 1:50, 1:250, and 1:1250 dilutions prepared in M199-CS medium containing 100 ng/ml FGF. The dilutions of B18 were added to the starved cells and incubated for 72 hr at 37°C. The cells were then radiolabeled by [<sup>3</sup>H]-thymidine for 6 hr. Radiolabeled cells were washed with PBS and trypsinized for liquid scintillation counting. Results were plotted using Kaleidograph software. The results are shown in Fig. 3. In the figure, "Med" is the mock control, "B18" and "B18-1" were conditioned medium from two independent transfections of COS with DNA encoding human CTLA-8 (B18). IFN $\gamma$  was used as a positive control angiostatic (i.e., angiogenesis inhibition) activity. These data demonstrate that human CTLA-8 (B18) inhibits angiogenesis.

#### Example 4

##### Hematopoietic Activity of Human CTLA-8

15 The hematopoietic activity of human CTLA-8 (B18) expressed *in vivo* was examined by construction of a recombinant adenovirus vector.

The B18 cDNA in the expression plasmid Adori 2-12 B18 was driven by the cytomegalovirus(CMV) immediate early promoter and enhancer.

20 The Adori 2-12 vector was created by addition of an SV40 origin and enhancer to a known adenovirus vector (Barr et al., Gene Therapy 1:51 (1994); Davidson et al., Nature Genetics 3:219 (1993)). The HindIII/BamHI fragment encoding the SV40 origin and enhancer was isolated from the pMT2 mammalian expression vector, blunted with Klenow and cloned into the NotI site (blunted with Klenow) of the Ad5 expression vector.

25 The vector was derived by digesting pNOT-B18 cDNA with *SalI*, filling in the 5' overhang with Klenow to generate a blunt end and digesting with *EcoRI* to isolate the B18 cDNA. The blunted- *EcoRI* B18 fragment was inserted into the restriction sites *EcoRV-EcoRI* of the adenovirus vector Adori 2-12. The CMV-B18 expression cassette was located downstream of the SV40 origin and enhancer, and 0-1 map units of the left hand end of the adenovirus type 5(Ad5). The SV40 splice donor and

30

acceptor were located between the CMV promoter and B18 cDNA. Following the insert was SV40 poly A site, 9-16 map units of Ad5 and the puc 19 origin.

A recombinant adenovirus was generated by homologous recombination in 293 cells. *AscI* linearized Adori 2-12 B18 and *ClaI* digested AdCMVlacZ were introduced  
5 into the 293 cells using lipofectamine. Recombinant adenovirus virus was isolated and amplified on 293 cells. The virus was released from infected 293 cells by three cycles of freeze-thawing. The virus was further purified by two cesium chloride centrifugation gradients and dialyzed against PBS 4°C. Following dialysis of the virus glycerol was added to a concentration of 10 % and the virus was stored at - 70 °C until  
10 use. The virus was characterized by expression of the transgene, plaque forming units on 293 cells, particles/ml and Southern analysis of the virus.

A single dose of  $5 \times 10^{10}$  particles of recombinant adenovirus encoding B18 was injected into the tail vein of male C57/bl6 mice, age 7-8 weeks. Control mice received an adenovirus encoding B-galactosidase. Four mice from each experimental  
15 group were killed on day 7 and 14. Blood was collected and automated hematologic analysis was performed using a Baker 9000. Differential counts were performed on blood smears. Tissue was harvested, fixed in formalin, and stained with hematoxylin and eosin for histopathology. In the first set of experiments, serum and tissues were analyzed 7 and 14 days post injection. A slight increase in peripheral platelet counts  
20 were observed. The animals that received B18 exhibited a slight increase in spleen size. Macroscopic analysis of the spleen showed an increase in splenic extramedullary hematopoiesis on day 7 compared to the control. These results showed a hematopoietic growth activity associated with B18.

In a second set of experiments  $5 \times 10^{10}$  particles of recombinant adenovirus encoding B18 were injected into the tail vein of male C57/bl6 mice, age 17-18 weeks.  
25 Control mice received an adenovirus encoding B-galactosidase. Blood samples were collected via retro-orbital sinus on days 2, 5, 7, 10, 14, and 21. The hematologic analyses were performed on the Baker 9000 automated cell counter with murine-specific settings. Analyses included WBC, RBC, HCT and PLT counts. Blood smears  
30 were prepared and stained with Wright-Geimsa for WBC differentials based on a 100 cellcount. Reticulocytes and reticulated platelets were quantitated using flow

cytometry. Four mice from each group were killed on days 7, 14, and 21. In addition to peripheral blood analysis, serum was collected via cardiac puncture for quantitation of systemic IL-6 using a commercial kit (Endogen). Spleen and liver were collected for histopathology, spleen and bone marrow hematopoietic progenitors were quantitated, and bone marrow smears were prepared and stained with Wright-Geimsa for cell counts.

Administration of adenovirus encoding B18 resulted in a marked increase in peripheral blood neutrophils and WBC (Fig. 4). Maximum increases in neutrophils were observed at day 5 and day 7. The control mice showed little difference at day 5 and day 7. Peripheral blood neutrophils were similar in the control mice and mice that received B18 at day 21. In both the B18 and control groups an increase in white blood cells was also observed. The mice that received B18 had a greater increase in WBC between day 2 and day 7. By Day 21 a more pronounced increase was observed in the B-gal group. No other changes in cellular chemistries were observed (Table I).

Bone marrow cellularity was calculated from pooled femurs in each group (Table III). No significant differences were observed in either group. No significant changes were observed in bone marrow hematopoietic progenitors from day 7, 14, and 21. The CFU-GM, BFU-E and CFU-MEG in the B18 mice were similar to the B-gal control (Table II).

Administration of the adenovirus encoding B18 resulted in an increase in CFU-GM (myeloid) and BFU-E (erythroid) progenitors in the spleen compared to animals that received the B-gal virus on day 7. The increase in progenitors in the B18 mice was 11-fold in CFU-GM and a 52-fold in BFU-E (Table II). There was a 2-fold increase in CFU-MEG at day 7 for the B18 mice. By day 21 no significant differences were observed in splenic CFU-MEG or BFU-E between the groups (Table II). A 3-fold decrease in CFU-GM was observed in mice that received adenovirus encoding B18. A slight increase in spleen size at day 7 was observed in the B18 group. This is consistent with an increase in splenic cellularity. By day 14 and day 21 spleen weights were similar to the control group (Table III). Macroscopic analysis of the spleen showed an increase in splenic extramedullary hematopoiesis of the B18 mice on day 7 compared to the control.

The bone marrow myeloid: erythroid ratios (Table IV) suggest a granulocytic hyperplasia with a possible erythroid hypoplasia in mice that received adenovirus B18 on day 7. By day 21 the ratio in the B-gal group was higher. No changes were observed in the IL6 serum levels.

- 5            These results show a hematopoietic activity associated with the administration of adenovirus encoding B18 (human CTLA-8). Increases in neutrophils and white blood cells were observed at day 7 in animals that received B18 adenovirus. The data showed that B18 resulted in increase in splenic CFU-GM and BFU-E 7 days post administration compared to the control animals. Splenic extramedullary
- 10           hematopoiesis on day 7 support that B18 exhibits a hematopoietic growth activity. These data suggest that B18 may mobilize early hematopoietic precursors.

Table I: Peripheral hematology for day 2, 5, 7, 10, 14, and 21.

Study AS4-43...B18 (Platelets) Day 2...4-25-96														
Group A	WBC x10 <sup>3</sup> /μL	Neutro %	ANC x10 <sup>3</sup> /μL	Lymphs %	ALC x10 <sup>3</sup> /μL	Eos %	Monos %	RBC x10 <sup>6</sup> /μL	Retics %	Abs Retics x10 <sup>6</sup> /μL	HCT %	PLT x10 <sup>3</sup> /μL	RPR %	Abs RPR x10 <sup>3</sup> /μL
B-Qul #1	5.4	40	2.18	54	2.92	0	6	10.88	3.65	0.40	48.0	836	11.84	99.82
B-Qul #2	7.4	25	1.85	65	4.81	3	7	12.34	2.04	0.25	56.6	900	10.10	90.90
B-Qul #3	6.6	40	2.72	52	3.54	2	6	11.28	3.28	0.37	51.8	894	9.77	87.34
B-Qul #4	8.8	23	2.02	84	5.63	1	12	12.00	2.55	0.31	54.8	840	10.63	89.29
AVG	7.1	32.0	2.19	59.8	4.22	1.5	7.8	11.82	2.88	0.33	52.8	868	10.81	91.84
SEM	0.7	4.6	0.19	3.4	0.51	0.5	1.4	0.33	0.36	0.03	1.9	17	0.49	2.76
B18 #1	11.4	59	6.73	31	3.53	1	8	11.18	4.88	0.54	52.4	1242	14.92	185.31
B18 #2	9.2	30	2.76	62	5.70	3	5	10.14	3.97	0.40	48.0	632	10.90	68.89
B18 #3	5.0	51	2.55	40	2.00	0	8	11.18	3.23	0.38	61.2	632	11.18	93.02
B18 #4	6.4	41	2.62	65	3.52	0	4	10.80	3.09	0.33	49.2	904	17.31	156.48
AVG	8.0	45.3	3.67	47.0	3.49	1.0	6.8	10.82	3.79	0.41	50.2	803	12.58	125.82
SEM	1.4	8.3	1.02	7.0	0.76	0.7	1.2	0.34	0.41	0.05	1.0	127	1.85	27.07

Study AS4-49...B18 (Platelets) Day 5...4-29-96														
Group B	WBC x10 <sup>3</sup> /μL	Neutro %	ANC x10 <sup>3</sup> /μL	Lymphs %	ALC x10 <sup>3</sup> /μL	Eos %	Monos %	RBC x10 <sup>6</sup> /μL	Retics %	Abs Retics x10 <sup>6</sup> /μL	HCT %	PLT x10 <sup>3</sup> /μL	RPR %	Abs RPR x10 <sup>3</sup> /μL
B-Qul #1	7.0	14	1.06	78	8.83	3	5	11.26	5.25	0.59	52.4	1082	15.31	167.62
B-Qul #2	10.6	20	2.12	78	8.27	1	1	10.72	4.71	0.50	49.4	994	17.37	172.06
B-Qul #3	8.8	18	1.51	69	6.07	2	11	11.12	3.40	0.38	51.2	918	9.55	87.48
B-Qul #4	10.8	38	4.10	58	6.26	0	1.5	10.22	6.21	0.63	47.0	1052	13.93	152.12
AVG	9.5	22.5	2.20	70.8	8.63	0.8	5.9	10.83	4.89	0.53	50.0	1021	14.09	145.02
SEM	0.8	6.3	0.67	4.8	0.55	0.4	2.1	0.25	0.59	0.06	1.2	41	1.57	19.87
B18 #1	14.8	18	2.86	71	10.51	1	10	12.66	2.31	0.29	57.2	1204	7.57	61.14
B18 #2	14.2	37	5.25	53	7.53	2	8	9.80	3.32	0.33	44.8	888	14.33	127.25
B18 #3	12.8	30	3.84	59	7.55	1	10	12.12	4.12	0.50	53.6	1134	10.15	115.10
B18 #4	16.0	58	9.28	37	5.92	0	8	11.04	3.93	0.43	50.8	1166	15.75	183.85
AVG	14.5	35.8	5.26	55.0	7.88	1.0	8.3	11.41	3.42	0.39	52.0	1098	11.95	128.28
SEM	0.7	8.4	1.44	7.1	0.96	0.4	1.2	0.63	0.41	0.05	2.8	71	1.85	18.81

Study AS4-49...B18 (Platelets) Day 7...4-30-96														
Group C	WBC x10 <sup>3</sup> /μL	Neutro %	ANC x10 <sup>3</sup> /μL	Lymphs %	ALC x10 <sup>3</sup> /μL	Eos %	Monos %	RBC x10 <sup>6</sup> /μL	Retics %	Abs Retics x10 <sup>6</sup> /μL	HCT %	PLT x10 <sup>3</sup> /μL	RPR %	Abs RPR x10 <sup>3</sup> /μL
B-Qul #1	13.2	14	2.19	69	10.49	1	18	11.04	3.54	0.39	50.8	862	12.45	107.41
B-Qul #2	14.0	12	1.88	81	11.34	0	7	11.36	5.05	0.57	52.8	1104	14.91	164.81
B-Qul #3	14.8	14	2.07	73	10.80	1	12	10.92	5.42	0.59	49.8	952	11.49	108.38
AVG	14.7	13.3	1.86	74.3	10.88	0.7	11.7	11.11	4.87	0.52	51.0	979	12.85	127.13
SEM	0.4	0.7	0.14	3.5	0.25	0.3	2.5	0.14	0.54	0.06	0.9	71	1.02	18.75
B18 #1	10.4	33	6.40	62	12.03	0	8	10.14	2.83	0.30	45.2	864	12.80	116.56
B18 #2	25.4	39	8.91	53	13.48	0	8	9.46	8.05	0.87	43.8	1288	12.49	190.67
B18 #3	23.6	44	10.38	50	11.80	0	6	9.74	5.17	0.50	44.4	1076	15.41	185.81
B18 #4	12.8	15	1.52	75	8.60	0	10	9.54	6.25	0.60	43.4	1136	15.68	180.40
AVG	20.3	32.8	7.15	60.0	11.72	0.0	7.3	9.72	5.10	0.49	44.2	1081	14.15	184.22
SEM	2.8	6.3	1.98	8.6	0.80	0.0	1.1	0.15	0.76	0.07	0.4	88	0.87	15.19

Study AS4-49...B18 (Platelets) Day 10...5-3-96														
Group A	WBC x10 <sup>3</sup> /μL	Neutro %	ANC x10 <sup>3</sup> /μL	Lymphs %	ALC x10 <sup>3</sup> /μL	Eos %	Monos %	RBC x10 <sup>6</sup> /μL	Retics %	Abs Retics x10 <sup>6</sup> /μL	HCT %	PLT x10 <sup>3</sup> /μL	RPR %	Abs RPR x10 <sup>3</sup> /μL
B-Qul #1	18.8	17	3.18	69	12.83	3	11	10.22	12.41	1.27	46.8	1460	16.20	236.52
B-Qul #2	13.2	18	2.11	79	10.43	1	4	10.48	6.00	0.63	46.8	1128	14.48	163.33
B-Qul #3	19.8	18	3.14	74	14.60	0	10	10.72	6.25	0.67	49.4	1338	18.58	221.84
B-Qul #4	18.6	21	3.91	72	13.39	3	4	10.44	7.59	0.79	48.4	1088	14.35	153.28
AVG	17.6	17.5	3.09	73.5	12.79	1.8	7.3	10.47	8.08	0.84	46.4	1249	18.40	183.74
SEM	1.5	1.3	0.37	2.1	0.88	0.8	1.9	0.10	1.49	0.16	0.8	91	0.58	20.78
B18 #1	14.2	33	4.69	56	7.95	5	8	8.70	11.97	1.04	39.2	1760	14.49	253.02
B18 #2	17.0	35	6.16	57	10.03	1	7	9.04	9.46	0.88	42.0	1104	18.68	208.44
B18 #3	16.2	39	6.32	57	8.23	1	3	4.74	16.77	0.79	22.4	894	20.19	290.06
B18 #4	14.2	25	3.55	68	9.37	1	8	9.30	9.83	0.92	42.0	1416	16.81	238.03
AVG	16.8	33.0	6.18	59.0	8.15	2.0	6.0	7.95	12.04	0.80	36.4	1294	19.84	240.81
SEM	0.8	2.9	0.68	2.3	0.43	1.0	1.1	1.08	1.87	0.05	4.7	189	3.24	11.77

Study AS4-49...B18 (Platelets) Day 14...5-7-96														
Group B	WBC x10 <sup>3</sup> /μL	Neutro %	ANC x10 <sup>3</sup> /μL	Lymphs %	ALC x10 <sup>3</sup> /μL	Eos %	Monos %	RBC x10 <sup>6</sup> /μL	Retics %	Abs Retics x10 <sup>6</sup> /μL	HCT %	PLT x10 <sup>3</sup> /μL	RPR %	Abs RPR x10 <sup>3</sup> /μL
B-Qul #1	17.8	18	3.20	74	13.17	0	8	10.06	5.57	0.63	50.8	1360	11.03	160.01
B-Qul #2	20.4	28	5.30	66	12.48	1	7	10.92	7.07	0.77	50.8	1618	8.18	132.19
B-Qul #3	16.0	7	1.12	90	14.40	1	3	11.36	6.41	0.73	52.8	1298	7.36	95.53
B-Qul #4	15.0	36	6.45	57	10.28	1	6	9.30	7.82	0.71	43.0	1672	10.05	168.04
AVG	18.1	21.4	4.03	71.8	12.82	0.8	6.0	10.81	6.77	0.71	49.4	1487	8.18	136.44
SEM	0.8	6.1	1.16	7.0	0.89	0.3	1.1	0.45	0.36	0.03	2.3	83	0.84	15.48
B18 #1	15.4	9	1.39	81	12.47	1	9	10.82	5.74	0.61	48.2	1282	9.51	120.52
B18 #2	15.4	31	4.77	58	8.83	2	9	9.76	10.33	1.01	44.8	1092	14.29	156.05
B18 #3	13.4	42	5.63	39	5.23	0	19	10.34	4.99	0.52	46.6	1378	15.79	217.27
B18 #4	11.8	57	6.61	34	3.94	2	7	9.38	8.57	0.52	43.0	1092	16.66	181.23
AVG	14.0	34.8	4.60	53.0	7.84	1.3	11.0	10.03	6.66	0.66	45.8	1206	14.06	168.82
SEM	0.9	10.1	1.18	10.7	1.93	0.8	2.7	0.28	1.23	0.12	1.1	79	1.69	20.84

Study AS4-49...B18 (Platelets) Day 21...5-14-96														
Group A	WBC x10 <sup>3</sup> /μL	Neutro %	ANC x10 <sup>3</sup> /μL	Lymphs %	ALC x10 <sup>3</sup> /μL	Eos %	Monos %	RBC x10 <sup>6</sup> /μL	Retics %	Abs Retics x10 <sup>6</sup> /μL	HCT %	PLT x10 <sup>3</sup> /μL	RPR %	Abs RPR x10 <sup>3</sup> /μL
B-Qul #1	23.4	23	5.84	67	17.02	0	10	9.22	6.15	0.75	42.8	1776	9.61	170.87
B-Qul #2	19.8	19	3.72	69	13.52	0	12	9.50	9.95	0.95	44.4	1682	9.44	156.89
B-Qul #3	27.8	11	3.04	82	22.63	3	4	9.74	8.84	0.88	43.8	1684	11.45	192.82
B-Qul #4	28.0	13	3.84	83	23.24	0	4	9.04	7.54	0.68	41.8	1346	10.48	141.06
AVG	25.2	16.5	4.08	75.3	19.10	0.8	7.5	9.38	8.62	0.81	43.7	1617	10.25	185.36
SEM	1.9	2.8	0.61	4.2	2.33	0.8	2.1	0.15	0.52	0.06	0.8	84	0.46	10.87
B18 #1	18.8	9	1.87	83	15.44	1	7	9.84	7.40	0.73	43.8	1642	8.37	137.44
B18 #2	18.2	48	7.78	45	7.29	1	6	11.10	7.87	0.88	51.4	1670	10.78	212.37
B18 #3	14.8	53	7.84	42	8.22	0	5	7.52	80.28	1.52	36.0	1568	10.50	164.84
B18 #4	15.2	18	2.74	74	11.25	2	6	9.64	7.22	0.68	44.1	1422	7.60	108.07
AVG	18.2	32.0	6.01	61.0	10.65	1.0	6.0	8.53	10.64	0.95	44.1	1651	9.31	155.83
SEM	0.8	10.9	1.83	10.3	2.10	0.4	0.4	0.74	3.21	0.20	2.8	116	0.78	22.18

Table II: Bone marrow and Splenic Hematopoietic Progenitors

	CFU-MEG		CFU-GM		BFU-E	
	B-Gal	B18	B-Gal	B18	B-Gal	B18
Bone Marrow*						
Day 7	16.0 ± 3.5	15.7 ± 3.1	307 ± 117	241 ± 78	51 ± 19	25 ± 11
Day 14	10.7 ± 2.3	15.3 ± 1.2	233 ± 15	373 ± 35	30 ± 10	60 ± 30
Day 21	5.7 ± 0.6	6.7 ± 3.1	170 ± 17	160 ± 27	40 ± 10	27 ± 6
Spleen**						
Day 7	9.3 ± 1.6	19.5 ± 1.5	27 ± 3	298 ± 6	1.3 ± 1.2	68 ± 10
Day 14	9.7 ± 0.6	12.7 ± 0.6	267 ± 32	197 ± 21	33 ± 6	10 ± 10
Day 21	17.0 ± 1.0	19.3 ± 2.5	187 ± 6	73 ± 15	23 ± 6	23 ± 6

Hematopoietic precursors were determined from pooled spleen and bone marrow samples from four animals in each group. For quantitation of CFU-GM and BFU-E, either  $1 \times 10^4$  bone marrow cells or  $1 \times 10^5$  spleen cells were added to complete alpha methylcellulose medium (0.9% methylcellulose in alpha medium, 30% fetal bovine serum, 1% bovine serum albumin,  $10^{-4}$ M 2-mercaptoethanol, 2 mM L-glutamine, 2% murine spleen cell conditioned medium, and 3 U/mL erythropoietin) and aliquoted into 35 mm tissue culture dishes in a final volume of 1.0 mL. Cultures were incubated for 7 days at 37°C, and 5% CO<sub>2</sub>. Microscopic colonies were defined as clusters of 50 or more cells. For quantitation of CFU-MEG, either  $1 \times 10^5$  bone marrow cells or  $1 \times 10^6$  spleen cells were added to complete alpha methylcellulose medium and incubated as described above. Megakaryocyte colonies were defined as a group of 3 or more cells.

\*Bone marrow progenitors are represented as mean ± sd number of colonies per  $10^5$  cells.

\*\*Spleen progenitors are represented as mean ± sd number of colonies per  $10^6$  cells.

Table III: Spleen Weights and Femur Cellularity

Spleen Wt. (Mg)	B-Gal	B18		Femur Cellularity ( $\times 10^6$ )	B-Gal	B18
Day 7	187 $\pm$ 19	224 $\pm$ 29		Day 7	28	23
Day 14	175 $\pm$ 13	170 $\pm$ 10		Day 14	28	27
Day 21	174 $\pm$ 21	151 $\pm$ 27		Day 21	28	26

Spleen weights were determined at time of sacrifice are represented as means  $\pm$  sd from four animals.

Table IV: Bone Marrow Myeloid:Erythroid Ratios

Group	Mouse #	Day 7	Day 14	Day 21
B-gal	1	1.43	2.12	5.78
	2	0.91	2.46	5.83
	3	1.62	1.03	3.66
	4		5.44	4.82
<b>AVG</b>		<b>1.32</b>	<b>2.76</b>	<b>5.02</b>
<b>SD</b>		<b>0.37</b>	<b>0.37</b>	<b>1.89</b>
B-gal	1	5.59	2.01	2.02
	2	6.51	1.25	2.13
	3	5.49	1.58	1.81
	4	0.50	2.51	2.92
<b>AVG</b>		<b>4.52</b>	<b>1.86</b>	<b>2.22</b>
<b>SD-</b>		<b>1.29</b>	<b>2.72</b>	<b>0.56</b>

All entries represent the number of myeloid cells per 1 erythroid cell. Normal mouse ratios are approximately 1:1 to 2:1.

#### Example 5

#### Additional Experiments Relating to

#### Hematopoietic Activity of Human CTLA-8

B18 (human CTLA-8) was tested for the ability to induce production of factors having hematopoietic activity in a factor-dependent cell proliferation assay using the

human erythroleukemic cell line, TF-1 (Kitamura et al., J. Cell Physiol. 140:323 (1989)). The cells were initially grown in the presence of rhGMCSF (100 U/ml). The cells were fed three days prior to setting up the assay. The assay conditions were as follows:

5	cells/well	5000/200 $\mu$ l
	incubation time	3 days
	pulse time	4 hours
	amount of tritiated thymidine	0.5 $\mu$ Ci/well
	counting time	1 minute
10	replicates	2

B18 alone, conditioned medium (CM) from B18 induced HS-5 cells were assayed. Buffer alone, CM from HS-5 cells induced with buffer and CM from uninduced HS-5 cells were assayed as controls. Results are shown in Fig. 5. B18 (human CTLA-8) demonstrated an ability to induce production of factors which induced TF-1 proliferation. This activity was substantially eliminated by the addition of anti-GMCSF antibodies. These data demonstrate that human CTLA-8 (B18) is able to induce hematopoiesis. Particularly, without being bound by any theory, it appears that human CTLA-8 (B18) induces production of GM-CSF and/or IL-3.

20

#### Example 6

##### Ability of Human CTLA-8 to Induce Production of IL-6 and IL-8

MRC5 cells were incubated in the presence of human CTLA-8 (B18) and production of IL-6 and IL-8 were measured. Herpes CTLA-8 (IL-17) was used as a positive control. Applicants' human CTLA-8 (B18) demonstrated titratable production of both IL-6 and IL-8 (see Figs. 6 and 7).

25

All patent and literature references cited herein are incorporated by reference as if fully set forth.



## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: Jacobs, Kenneth  
Kelleher, Kerry  
Carlin, McKeough  
Goldman, Samuel  
Pittman, Debra  
Mi, Sha  
Neben, Steven  
Giannotti, JoAnn  
Golden Fleet, Margaret
- (ii) TITLE OF INVENTION: Human CTLA-8 and Uses of CTLA-8-Related Proteins
- (iii) NUMBER OF SEQUENCES: 9
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Genetics Institute, Inc.
  - (B) STREET: 87 CambridgePark Drive
  - (C) CITY: Cambridge
  - (D) STATE: Massachusetts
  - (E) COUNTRY: USA
  - (F) ZIP: 02140
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER:
  - (B) FILING DATE:
  - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: Brown, Scott A.
  - (B) REGISTRATION NUMBER: 32,724
  - (C) REFERENCE/DOCKET NUMBER: GI5262
- (ix) TELECOMMUNICATION INFORMATION:
  - (A) TELEPHONE: (617) 498-8224
  - (B) TELEFAX: (617) 876-5851

## (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 813 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO
- (ix) FEATURE:
  - (A) NAME/KEY: CDS
  - (B) LOCATION: 56..544

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GGGAAGATAC ATTACAGAA AGAGCTTCCT GCACAAAGTA AGCCACCAGC GCAAC ATG	58
Met	
1	
ACA GTG AAG ACC CTG CAT GGC CCA GCC ATG GTC AAG TAC TTG CTG CTG	106
Thr Val Lys Thr Leu His Gly Pro Ala Met Val Lys Tyr Leu Leu Leu	
5 10 15	
TCG ATA TTG GGG CTT GCC TTT CTG AGT GAG GCG GCA GCT CGG AAA ATC	154
Ser Ile Leu Gly Leu Ala Phe Leu Ser Glu Ala Ala Ala Arg Lys Ile	
20 25 30	
CCC AAA GTA GGA CAT ACT TTT TTC CAA AAG CCT GAG AGT TGC CCG CCT	202
Pro Lys Val Gly His Thr Phe Phe Gln Lys Pro Glu Ser Cys Pro Pro	
35 40 45	
GTG CCA GGA GGT AGT ATG AAG CTT GAC ATT GGC ATC ATC AAT GAA AAC	250
Val Pro Gly Gly Ser Met Lys Leu Asp Ile Gly Ile Ile Asn Glu Asn	
50 55 60 65	
CAG CGC GTT TCC ATG TCA CGT AAC ATC GAG AGC CGC TCC ACC TCC CCC	298
Gln Arg Val Ser Met Ser Arg Asn Ile Glu Ser Arg Ser Thr Ser Pro	
70 75 80	
TGG AAT TAC ACT GTC ACT TGG GAC CCC AAC CGG TAC CCC TCG GAA GTT	346
Trp Asn Tyr Thr Val Thr Trp Asp Pro Asn Arg Tyr Pro Ser Glu Val	
85 90 95	
GTA CAG GCC CAG TGT AGG AAC TTG GGC TGC ATC AAT GCT CAA GGA AAG	394
Val Gln Ala Gln Cys Arg Asn Leu Gly Cys Ile Asn Ala Gln Gly Lys	
100 105 110	
GAA GAC ATC TCC ATG AAT TCC GTT CCC ATC CAG CAA GAG ACC CTG GTC	442
Glu Asp Ile Ser Met Asn Ser Val Pro Ile Gln Gln Glu Thr Leu Val	
115 120 125	
GTC CGG AGG AAG CAC CAA GGC TGC TCT GTT TCT TTC CAG TTG GAG AAG	490
Val Arg Arg Lys His Gln Gly Cys Ser Val Ser Phe Gln Leu Glu Lys	
130 135 140 145	
GTG CTG GTG ACT GTT GGC TGC ACC TGC GTC ACC CCT GTC ATC CAC CAT	538
Val Leu Val Thr Val Gly Cys Thr Cys Val Thr Pro Val Ile His His	
150 155 160	
GTG CAG TAAGAGGTGC ATATCCACTC AGCTGAAGAA GCTGTAGAAA TGCCACTCCT	594
Val Gln	
TACCCAGTGC TCTGCAACAA GTCCTGTCTG ACCCCCAATT CCCTCCACTT CACAGGACTC	654
TTAATAAGAC CTGCACGGAT GGAAACAGAA AATATTCACA ATGTATGTGT GTATGTACTA	714
CACCTTTATAT TTGATATCTA AAATGTTAGG AGAAAAATTA ATATATTCAG TGCTAATATA	774
ATAAAGTATT AATAATTAA AAATAAAAAA AAAAAAAA	813

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 163 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Thr Val Lys Thr Leu His Gly Pro Ala Met Val Lys Tyr Leu Leu  
 1 5 10 15  
 Leu Ser Ile Leu Gly Leu Ala Phe Leu Ser Glu Ala Ala Ala Arg Lys  
 20 25 30  
 Ile Pro Lys Val Gly His Thr Phe Phe Gln Lys Pro Glu Ser Cys Pro  
 35 40 45  
 Pro Val Pro Gly Gly Ser Met Lys Leu Asp Ile Gly Ile Ile Asn Glu  
 50 55 60  
 Asn Gln Arg Val Ser Met Ser Arg Asn Ile Glu Ser Arg Ser Thr Ser  
 65 70 75 80  
 Pro Trp Asn Tyr Thr Val Thr Trp Asp Pro Asn Arg Tyr Pro Ser Glu  
 85 90 95  
 Val Val Gln Ala Gln Cys Arg Asn Leu Gly Cys Ile Asn Ala Gln Gly  
 100 105 110  
 Lys Glu Asp Ile Ser Met Asn Ser Val Pro Ile Gln Gln Glu Thr Leu  
 115 120 125  
 Val Val Arg Arg Lys His Gln Gly Cys Ser Val Ser Phe Gln Leu Glu  
 130 135 140  
 Lys Val Leu Val Thr Val Gly Cys Thr Cys Val Thr Pro Val Ile His  
 145 150 155 160  
 His Val Gln

## (2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 461 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

- (ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 6..455

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CCACC ATG TGC CTG ATG CTG TTG CTG CTA CTG AAC CTG GAG GCT ACA 47  
 Met Cys Leu Met Leu Leu Leu Leu Leu Leu Glu Ala Thr  
 1 5 10  
 GTG AAG GCA GCG GTA CTC ATC CCT CAA AGT TCA GTG TGT CCA AAC GCC 95  
 Val Lys Ala Ala Val Leu Ile Pro Gln Ser Ser Val Cys Pro Asn Ala  
 15 20 25 30  
 GAG GCC AAT AAC TTT CTC CAG AAC GTG AAG GTC AAC CTG AAA GTC ATC 143  
 Glu Ala Asn Asn Phe Leu Gln Asn Val Lys Val Asn Leu Lys Val Ile  
 35 40 45

AAC TCC CTT AGC TCA AAA GCG AGC TCG AGA AGG CCC TCA GAT TAC CTC	191
Asn Ser Leu Ser Ser Lys Ala Ser Ser Arg Arg Pro Ser Asp Tyr Leu	
50 55 60	
AAC CGT TCC ACT TCA CCC TGG ACT CTG AGC CGC AAT GAG GAC CCT GAT	239
Asn Arg Ser Thr Ser Pro Trp Thr Leu Ser Arg Asn Glu Asp Pro Asp	
65 70 75	
AGA TAT CCT TCT GTG ATC TGG GAG GCA CAG TGC CGC CAC CAG CGC TGT	287
Arg Tyr Pro Ser Val Ile Trp Glu Ala Gln Cys Arg His Gln Arg Cys	
80 85 90	
GTC AAC GCT GAG GGG AAG TTG GAC CAC CAC ATG AAT TCT GTT CTC ATC	335
Val Asn Ala Glu Gly Lys Leu Asp His His Met Asn Ser Val Leu Ile	
95 100 105 110	
CAG CAA GAG ATA CTA GTC CTG AAG AGG GAG CCT GAG AAG TGC CCC TTC	383
Gln Gln Glu Ile Leu Val Leu Lys Arg Glu Pro Glu Lys Cys Pro Phe	
115 120 125	
ACT TTC CGG GTG GAG AAG ATG CTG GTG GGC GTG GGC TGC ACC TGC GTT	431
Thr Phe Arg Val Glu Lys Met Leu Val Gly Val Gly Cys Thr Cys Val	
130 135 140	
TCC TCT ATT GTC CGC CAT GCG TCC TAATAA	461
Ser Ser Ile Val Arg His Ala Ser	
145 150	

## (2) INFORMATION FOR SEQ ID NO:4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 150 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Cys Leu Met Leu Leu Leu Leu Asn Leu Glu Ala Thr Val Lys	
1 5 10 15	
Ala Ala Val Leu Ile Pro Gln Ser Ser Val Cys Pro Asn Ala Glu Ala	
20 25 30	
Asn Asn Phe Leu Gln Asn Val Lys Val Asn Leu Lys Val Ile Asn Ser	
35 40 45	
Leu Ser Ser Lys Ala Ser Ser Arg Arg Pro Ser Asp Tyr Leu Asn Arg	
50 55 60	
Ser Thr Ser Pro Trp Thr Leu Ser Arg Asn Glu Asp Pro Asp Arg Tyr	
65 70 75 80	
Pro Ser Val Ile Trp Glu Ala Gln Cys Arg His Gln Arg Cys Val Asn	
85 90 95	
Ala Glu Gly Lys Leu Asp His His Met Asn Ser Val Leu Ile Gln Gln	
100 105 110	
Glu Ile Leu Val Leu Lys Arg Glu Pro Glu Lys Cys Pro Phe Thr Phe	
115 120 125	
Arg Val Glu Lys Met Leu Val Gly Val Gly Cys Thr Cys Val Ser Ser	
130 135 140	

Ile Val Arg His Ala Ser  
145 150

## (2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 459 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

- (ix) FEATURE:  
 (A) NAME/KEY: CDS  
 (B) LOCATION: 1..453

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

ATG ACA TTT AGA ATG ACT TCA CTT GTG TTA CTT CTG CTG CTG AGC ATA	48
Met Thr Phe Arg Met Thr Ser Leu Val Leu Leu Leu Leu Ser Ile	
1 5 10 15	
GAT TGT ATA GTA AAG TCA GAA ATA ACT AGT GCA CAA ACC CCA AGA TGC	96
Asp Cys Ile Val Lys Ser Glu Ile Thr Ser Ala Gln Thr Pro Arg Cys	
20 25 30	
TTA GCT GCT AAC AAT AGC TTT CCA CGG TCT GTG ATG GTT ACT TTG AGC	144
Leu Ala Ala Asn Asn Ser Phe Pro Arg Ser Val Met Val Thr Leu Ser	
35 40 45	
ATC CGT AAC TGG AAT ACC AGT TCT AAA AGG GCT TCA GAC TAC TAC AAT	192
Ile Arg Asn Trp Asn Thr Ser Ser Lys Arg Ala Ser Asp Tyr Tyr Asn	
50 55 60	
AGA TCT ACG TCT CCT TGG ACT CTC CAT CGC AAT GAA GAT CAA GAT AGA	240
Arg Ser Thr Ser Pro Trp Thr Leu His Arg Asn Glu Asp Gln Asp Arg	
65 70 75 80	
TAT CCC TCT GTG ATT TGG GAA GCA AAG TGT CGC TAC TTA GGA TGT GTT	288
Tyr Pro Ser Val Ile Trp Glu Ala Lys Cys Arg Tyr Leu Gly Cys Val	
85 90 95	
AAT GCT GAT GGG AAT GTA GAC TAC CAC ATG AAC TCA GTC CCT ATC CAA	336
Asn Ala Asp Gly Asn Val Asp Tyr His Met Asn Ser Val Pro Ile Gln	
100 105 110	
CAA GAG ATT CTA GTG GTG CGC AAA GGG CAT CAA CCC TGC CCT AAT TCA	384
Gln Glu Ile Leu Val Val Arg Lys Gly His Gln Pro Cys Pro Asn Ser	
115 120 125	
TTT AGG CTA GAG AAG ATG CTA GTG ACT GTA GGC TGC ACA TGC GTT ACT	432
Phe Arg Leu Glu Lys Met Leu Val Thr Val Gly Cys Thr Cys Val Thr	
130 135 140	
CCC ATT GTT CAC AAT GTA GAC TAAAAG	459
Pro Ile Val His Asn Val Asp	
145 150	

## (2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 151 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

```

Met Thr Phe Arg Met Thr Ser Leu Val Leu Leu Leu Leu Ser Ile
 1           5           10           15
Asp Cys Ile Val Lys Ser Glu Ile Thr Ser Ala Gln Thr Pro Arg Cys
          20           25           30
Leu Ala Ala Asn Asn Ser Phe Pro Arg Ser Val Met Val Thr Leu Ser
          35           40           45
Ile Arg Asn Trp Asn Thr Ser Ser Lys Arg Ala Ser Asp Tyr Tyr Asn
          50           55           60
Arg Ser Thr Ser Pro Trp Thr Leu His Arg Asn Glu Asp Gln Asp Arg
          65           70           75           80
Tyr Pro Ser Val Ile Trp Glu Ala Lys Cys Arg Tyr Leu Gly Cys Val
          85           90           95
Asn Ala Asp Gly Asn Val Asp Tyr His Met Asn Ser Val Pro Ile Gln
          100          105          110
Gln Glu Ile Leu Val Val Arg Lys Gly His Gln Pro Cys Pro Asn Ser
          115          120          125
Phe Arg Leu Glu Lys Met Leu Val Thr Val Gly Cys Thr Cys Val Thr
          130          135          140
Pro Ile Val His Asn Val Asp
          145          150
  
```

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 28 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CACAGGCATA CACAGGAAGA TACATTCA

28

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 25 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

TCTTGCTGGA TGGGAACGGA ATTCA

25

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 28 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ATACATTCAC AGAAGAGCTT CCTGCACA

28

5      What is claimed is:

1.      An isolated polynucleotide comprising a nucleotide sequence selected from the group consisting of:

         (a)      the nucleotide sequence of SEQ ID NO:1 from nucleotide 146 to nucleotide 544;

10           (b)      a nucleotide sequence capable of hybridizing to a nucleic acid sequence specified in (a);

         (c)      a nucleotide sequence varying from the sequence of the nucleotide sequence specified in (a) as a result of degeneracy of the genetic code; and

         (d)      an allelic variant of the nucleotide sequence specified in (a).

15

2.      The polynucleotide of claim 1 wherein said nucleotide sequence encodes a protein having CTLA-8 activity.

20      3.      The polynucleotide of claim 1 wherein said nucleotide sequence is operably linked to an expression control sequence.

4.      The polynucleotide of claim 1 comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 55 to nucleotide 544.

25      5.      A host cell transformed with the polynucleotide of claim 3.

6.      The host cell of claim 5, wherein said cell is a mammalian cell.

7.      A process for producing a human CTLA-8 protein, said process comprising:

30           (a)      growing a culture of the host cell of claim 5 in a suitable culture medium;  
         and

         (b)      purifying the human CTLA-8 protein from the culture.



- 5        8.        An isolated human CTLA-8 protein comprising an amino acid sequence selected from the group consisting of:
- (a)        the amino acid sequence of SEQ ID NO:2;
  - (b)        the amino acid sequence of SEQ ID NO:2 from amino acids 11 to 163;
  - (c)        the amino acid sequence of SEQ ID NO:2 from amino acids 29 to 163;
  - 10        (d)        the amino acid sequence of SEQ ID NO:2 from amino acids 31 to 163; and
  - (e)        fragments of (a), (b), (c) or (d) having CTLA-8 activity.
9.        The protein of claim 8 comprising the amino acid sequence of SEQ ID NO:2.
- 15        10.        The protein of claim 8 comprising the sequence from amino acid 29 to 163 of SEQ ID NO:2.
11.        A pharmaceutical composition comprising a human CTLA-8 protein of claim 8 and a pharmaceutically acceptable carrier.
- 20        12.        A human CTLA-8 protein produced according to the process of claim 7.
13.        A composition comprising an antibody which specifically reacts with a human CTLA-8 protein of claim 8.
- 25        14.        A method of treating a mammalian subject comprising administering a therapeutically effective amount of a composition of claim 11.
15.        A method of treating a mammalian subject comprising administering a therapeutically effective amount of a composition comprising a pharmaceutically acceptable carrier and a protein comprising an amino acid sequence selected from the group consisting of:
- 30        (a)        the amino acid sequence of SEQ ID NO:4;

- 5           (b)    the amino acid sequence of SEQ ID NO:4 from amino acids 18 to 150; and  
          (c)    fragments of (a) or (b) having CTLA-8 activity.

16.    The method of claim 15 wherein said protein comprises the amino acid sequence of SEQ ID NO:4.

10

17.    The method of claim 15 wherein said protein comprises the amino acid sequence of SEQ ID NO:4 from amino acids 18 to 150.

18.    A method of treating a mammalian subject administering a therapeutically effective amount of a composition comprising a pharmaceutically acceptable carrier and a protein comprising an amino acid sequence selected from the group consisting of:

15

- (a)    the amino acid sequence of SEQ ID NO:6;  
          (b)    the amino acid sequence of SEQ ID NO:6 from amino acids 19 to 151; and  
          (c)    fragments of (a) or (b) having CTLA-8 activity.

20

19.    The method of claim 18 wherein said protein comprises the amino acid sequence of SEQ ID NO:6.

20.    The method of claim 18 wherein said protein comprises the amino acid sequence of SEQ ID NO:6 from amino acids 19 to 151.

25

21.    The polynucleotide of claim 1 comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 86 to nucleotide 544.

30

22.    The protein of claim 8 comprising the sequence from amino acid 11 to 163 of SEQ ID NO:2.

- 5 23. A method of treating a mammalian subject comprising administering a therapeutically effective amount of a composition comprising a pharmaceutically acceptable carrier and IL-17 or an active fragment thereof.
- 10 24. The method of claim 14, 15, 18 or 23 wherein said subject is treated to produce an effect selected from the group consisting of inhibition of angiogenesis, inhibition of growth or proliferation of vascular endothelial cells, inhibition of tumor growth, inhibition of angiogenesis-dependent tissue growth, proliferation of myeloid cells or progenitors, proliferation of erythroid cells or progenitors, proliferation of lymphoid cells or progenitors, induction of IFN $\gamma$  production, induction of IL-3 production and induction of GM-CSF  
15 production.
25. The composition of claim 3 wherein said polynucleotide is contained in a vector suitable for *in vivo* expression in a mammalian subject.
- 20 26. The polynucleotide of claim 1 comprising the nucleotide sequence of SEQ ID NO:1 from nucleotide 139 to nucleotide 544.
27. The protein of claim 8 comprising the sequence from amino acid 31 to 163 of SEQ ID NO:2.



Figure 2



200K —  
97K —  
69K —  
46K —  
30K —  
23K —  
14.3K —



B18 M B18 M  
R R NR NR

Supernatant Cell Lysate

27




Fig. 3

3/7

Fig. 4A




Fig. 4B




Fig. 4C





Fig. 4D






Fig. 5

5/7







Fig. 6



# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 96/11889

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/19 C12N15/12 C12N15/24 C07K14/52 C07K14/725  
C07K14/54 A61K38/17 A61K38/19 A61K38/20 C12N5/10  
C07K16/24 C07K16/28 A61K48/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 95 18826 A (SCHERING CORP ; INST NAT SANTE RECH MED (FR)) 13 July 1995 cited in the application	1-3,5-7, 12, 15-20, 23-25
A	see page 3, line 21 - page 4, line 11 see page 12, line 1 - page 16, line 6 see page 30, line 1-13 Seq.ID:2 see page 53 Seq.ID:4 see page 55 Seq.ID:5 see page 56 Seq.ID:7/8 see page 58 - page 59 --- -/--	11,13,14

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&\* document member of the same patent family

Date of the actual completion of the international search

21 July 1997

Date of mailing of the international search report

25. 07. 97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+ 31-70) 340-3016

Authorized officer

Macchia, G

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 96/11889

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>THE JOURNAL OF IMMUNOLOGY, vol. 150, no. 12, 15 June 1993, pages 5445-5456, XP002035505 ROUVIER E. ET AL.: "CTLA-8, cloned from an activated T cell, bearing AU-rich messenger RNA instability sequences, and homologous to a Herpesvirus Saimiri gene" cited in the application see page 5445 - page 5446, line 11 see page 5449, right-hand column, line 26 - page 5450 see page 5451; figure 3 see page 5453, left-hand column, line 10 - right-hand column, line 16 ---</p>	15-20
X	<p>JOURNAL OF VIROLOGY, vol. 66, no. 8, August 1992, pages 5047-5058, XP000615399 ALBRECHT J -C ET AL: "PRIMARY STRUCTURE OF THE HERPESVIRUS SAIMIRI GENOME" Seq.ID:1 from nt.552 to nt.217 (reverse orientation) is 61.4% homologous to X64346 from nt.26931 to nt.27266. see page 5048, right-hand column, line 15-18 ORF13 see page 5049; table 1 ---</p>	1
A	<p>Seq.ID:1 from nt.552 to nt.217 (reverse orientation) is 61.4% homologous to X64346 from nt.26931 to nt.27266. see page 5048, right-hand column, line 15-18 ORF13 see page 5049; table 1 ---</p>	18-20
E	<p>WO 97 07198 A (GENETICS INST) 27 February 1997 see page 8, line 15-31 see page 9, line 26 - page 11, line 7 see page 13, line 16-27 see page 18, line 10 - page 19, line 25 see page 25, line 11-15 see page 27, line 5-17 see page 31, line 27-34 Seq.ID:11-12 see page 49 - page 51 ---</p>	1-14,21, 22,24-27
P,X	<p>JOURNAL OF IMMUNOLOGY, vol. 155, no. 12, 15 December 1995, pages 5483-5486, XP000602481 YAO Z ET AL: "HUMAN IL-17: A NOVEL CYTOKINE DERIVED FROM T CELLS" see page 5483 see page 5484; figure 1 ---</p>	1-3,5-7, 23
	-/--	

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 96/11889

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>THE JOURNAL OF EXPERIMENTAL MEDICINE, vol. 183, no. 6, 1 June 1996, pages 2593-2603, XP002035506 FOSSIEZ F. ET AL.: "T cell interleukin-17 induces stromal cells to produce proinflammatory and hematopoietic cytokines" see abstract see page 2594; figure 1 ---</p>	<p>1-3,5-7, 15-20, 23,24</p>
P,X	<p>IMMUNITY, vol. 3, no. 6, 1 December 1995, pages 811-821, XP000579309 YAO Z ET AL: "HERPESVIRUS SAIMIRI ENCODES A NEW CYTOKINE, IL-17, WHICH BINDS TO A NOVEL CYTOKINE RECEPTOR" see page 811 see page 815, left-hand column, line 50 - page 818, left-hand column ---</p>	<p>15-20, 23,24</p>
E	<p>WO 96 29408 A (IMMUNEX CORP) 26 September 1996 see page 1, line 1 - page 2, line 33 Seq.ID:8 see page 36 ---</p>	<p>18-20</p>
P,X	<p>GENE, vol. 168, no. 2, 12 February 1996, pages 223-225, XP002035631 YAO ET AL.: "Complete nucleotide sequence of the mouse CTLA8 gene" see page 225; figure 2 -----</p>	<p>1,2</p>

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 96/ 11889

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 14-20, 23, 24  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 14-20, 23, 24 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see continuation-sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

## INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 96/ 11889

FURTHER INFORMATION CONTINUED FROM PCT/ISA/210

- 1) claims 1-14, 21, 22, 25-27 all totally; claim 24 partially.

Isolated polynucleotide comprising Seq.ID:1, homologue sequences and derivatives. Vectors and transformed host cells. Process for producing human recombinant CTLA-8 protein. Isolated human CTLA-8 protein as in Seq.ID:2 and fragments. Pharmaceutical compositions and uses in therapy. Antibodies.

- 2) claims 15-17 all totally; claim 24 partially.

Therapeutical uses of the protein as in Seq.ID:4 or fragments.

- 3) claims 18-20 all totally; claim 24 partially.

Therapeutical uses of the protein as in Seq.ID:6 or fragments.

- 4) claims 23 all totally; claim 24 partially.

Therapeutical uses of IL-17 or active fragments.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 96/11889

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9518826 A	13-07-95	AU 1520895 A EP 0733069 A JP 9501572 T	01-08-95 25-09-96 18-02-97
WO 9707198 A	27-02-97	AU 6712396 A AU 6768596 A WO 9704097 A	18-02-97 12-03-97 06-02-97
WO 9629408 A	26-09-96	AU 5526396 A	08-10-96